❖ **Karyokinesis:**

- Involves division of nucleus.
- Nucleus develops a constriction at the center and becomes dumbell shaped.
- Constriction deepens and divides the nucleus into two.

❖ **Cytokinesis:**

- Involves division of cytoplasm.
- Plasma membrane develops a constriction along nuclear constriction.
- It deepens centripetally and finally divides the cell into two cells.

Example: Cells of mammalian cartilage, macronucleus of *Paramecium* and old degenerating cells of higher plants.

Drawbacks of Amitosis

- Causes unequal distribution of chromosomes.
- Can lead to abnormalities in metabolism and reproduction.

7.3.2 Mitosis

Mitosis occurs in shoot and root tips and other meristematic tissues of plants associated with growth. The number of chromosomes in the parent and the daughter (Progeny) cells remain the same so it is also called as **equational division**.

7.3.3 Closed and Open Mitosis

In closed mitosis, the nuclear envelope remains intact and chromosomes migrate to opposite poles of a spindle within the nucleus (Figure 7.3).

Example: Many single celled eukaryotes including yeast and slime molds.

In open mitosis, the nuclear envelope breaks down and then reforms around the 2 sets of separated chromosome.

Example: Most plants and animals

- Some animals are able to regenerate whole parts of the body.




Figure 7.3: Closed and Open mitosis

Mitosis is divided into four stages prophase, metaphase, anaphase and telophase (Figure 7.6).

Prophase

Prophase is the longest phase in mitosis. Chromosomes become visible as long thin thread like structure, condenses to form compact mitotic chromosomes. In plant cells initiation of spindle fibres takes place, nucleolus disappears. Nuclear envelope breaks down. Golgi apparatus and endoplasmic reticulum disappear.




Figure 7.4: Centromere

In animal cell the centrioles extend a radial array of microtubules (Figure 7.4) and reach the poles of the cell. This arrangement of microtubules is called **an aster**. Plant cells do not form asters.

Metaphase

Chromosomes (two sister chromatids) are attached to the spindle fibres by kinetochore of the centromere. The spindle fibres are made up of tubulin. The alignment of chromosome

XIth_Botany_Chapter_7_EM.indd 169 5/17/2020 9:12:09 PM