
there becomes new cell wall. The first stage of cell wall construction is a line dividing the newly forming cells called a **cell plate**. The cell plate eventually stretches right across the cell forming the middle lamella. Cellulose builds up on each side of the middle lamella to form the cell walls of two new plant cells.

Activity

Squash preparation of onion root tip to visualize and study various stages of mitosis.



A Culture of animal cells in which the cell cycles were asynchronous was incubated with 3H-Thymidine for 10 minutes. Autoradiography showed that 50% of the cells were labelled. If the cell cycle time (generation time) was 16 hrs how long was the S period?

Length of the S period = Fraction of cells in DNA replication × generation time

Length of the S period = 0.5 × 16 hours = 8 hours



Skin cells and the cells lining our gut are constantly dying and are being replaced by identical cells.

7.3.5 Significance of Mitosis

Exact copy of the parent cell is produced by mitosis (genetically identical).

1. **Genetic stability** – daughter cells are genetically identical to parent cells.
2. **Growth** – as multicellular organisms grow, the number of cells making up their tissue increases. The new cells must be identical to the existing ones.
3. **Repair of tissues** - damaged cells must be replaced by identical new cells by mitosis.
4. **Asexual reproduction** – asexual reproduction results in offspring that are identical to the parent. Example Yeast and Amoeba.
5. In flowering plants, structure such as bulbs, corms, tubers, rhizomes and runners are produced by mitotic division. When they separate from the parent, they form a new individual.
   The production of large numbers of offsprings in a short period of time, is possible only by mitosis. In genetic engineering and biotechnology, tissues are grown by mitosis (i.e. in tissue culture).
6. **Regeneration** – Arms of star fish.

7.3.6 Meiosis

In Greek *meioum* means to reduce. Meiosis is unique because of synapsis, homologous recombination and reduction division. Meiosis takes place in the reproductive organs. It results in the formation of gametes with half the normal chromosome number.

Haploid sperms are made in testes; haploid eggs are made in ovaries of animals.

In flowering plants meiosis occurs during microsporogenesis in anthers and megasporogenesis in ovule. In contrast to mitosis, meiosis produces cells that are not genetically identical. So meiosis has a key role in producing new genetic types which results in genetic variation.

Stages in Meiosis

Meiosis can be studied under two divisions i.e., meiosis I and meiosis II. As with mitosis, the cell is said to be in interphase when it is not dividing.

Prophase I is the longest and most complex stage in meiosis. Pairing of homologous chromosomes (bivalents) take place.

Meiosis I-Reduction Division

Prophase I – Prophase I is of longer duration and it is divided into 5 substages – Leptotene, Zygotene, Pachytene, Diplotene and Diakinesis (Figure 7.7).

XIth_Botany_Chapter_7_EM.indd 171 5/17/2020 9:12:11 PM