8.5 Proteins

Proteins are the most diverse of all macromolecule. Proteins make up 2/3 of total dry mass of a cell. The term protein was coined by **Gerardus Johannes Mulder** and is derived form a *greek* word proteos which means of the first rank.

Amino acids are building blocks of proteins. There are about 20 different amino acids exist naturally. All amino acids have a basic skeleton consisting of a carbon (a-carbon) linked to a basic amino group.




Figure 8.9: Structure of basic amino acid

(NH_2), an acidic carboxylic group (COOH) and a hydrogen atom (H) and side chain or variable R group. The amino acid is both an acid and a base and hence is called **amphoteric**.

A **zwitterion** also called as **dipolar ion**, is a molecule with two or more functional groups, of which at least one has a positive and other has a negative electrical charge and the net charge of the entire molecule is zero. The pH at which this happens is known as the **isoelectric point** (Figure 8.10).




Figure 8.10: Structure of amino acid

8.5.1 Classification of Amino acids

Based on the R group amino acids are classified as acidic, basic, polar, non-polar.

The amino group of one amino acid reacts with carboxyl group of other amino acid, forming a **peptide bond**. Two amino acids can react together with the loss of water to form a **dipeptide**. Long strings of amino acids linked by peptide bonds are called **polypeptides**. In 1953, Fred Sanger first sequenced the Insulin protein (Figure 8.11 a and b).





Figure 8.11(a): Formation of peptide bond



Linus Pauling and Robert Corey in 1951 proposed the α-helix and β sheet secondary structures of proteins. They were awarded Nobel prize in 1954



First protein Insulin was sequenced by Fred Sange.

XIth_Botany_Chapter_8_EM.indd 183 5/17/2020 9:14:47 PM