Disulfide Bond

Some amino acids like cysteine and methionine have sulphur. These form disulphide bridge between sulphur atoms and amino acids.

Hydrophobic Bond

This bond helps some protein to maintain structure. When globular proteins are in solution, their hydrophobic groups point inwards away from water.

8.5.5 Test for Proteins

The biuret test is used as an indicator for presence of protein as it gives a purple colour in the presence of peptide bonds (–C–N–). To protein solution, an equal quantity of sodium hydroxide solution is added and mixed. Then a few drops of 0.5% copper (II) sulphate is added with gentle mixing. A distinct purple colour develops without heating (Figure 8.15 a and b).




Figure 8.15(a): Biuret test




Figure 8.15(b): Colour intensity increases with increase in concentration



The more the distance between the sulphur atoms, the more the proteins bend; the more the hair curls.



8.6 Enzymes

Enzymes are globular proteins that catalyse the many thousands of metabolic reactions taking place within cells and organism. The molecules involved in such reactions are metabolites. Metabolism consists of chains and cycles of enzyme-catalysed reactions, such as respiration, photosynthesis, protein synthesis and other pathways. These reactions are classified as

- **anabolic** (building up of organic molecules). Synthesis of proteins from amino acids and synthesis of polysaccharides from simple sugars are examples of anabolic reactions.
- **catabolic** (breaking down of larger molecules). Digestion of complex foods and the breaking down of sugar in respiration are examples of catabolic reactions (Figure 8.16).




Figure 8.16: Enzyme reaction

Enzymes can be **extracellular enzyme** as secreted and work externally exported from cells. Eg. digestive enzymes; or **intracellular enzymes** that remain within cells and work there. These are found inside organelles or within cells. Eg. insulin.

XIth_Botany_Chapter_8_EM.indd 186 5/17/2020 9:14:50 PM