DO YOU KNOW?

ATP is not only higher energy compound present in a cell. There are other higher energy compounds also present. Example GTP (Guanosine Tri Phosphate) and UTP (Uridine Tri Phosphate).

14.5 Stages of Respiration

1. Glycolysis-conversion of glucose into pyruvic acid in cytoplasm of cell.
2. Link reaction-conversion of pyruvic acid into acetyl coenzyme-A in mitochondrial matrix.
3. Krebs cycle-conversion of acetyl coenzyme A into carbon dioxide and water in the mitochondrial matrix.
4. Electron transport chain to tranfer electrons remove hydrogen ions and tranfer electrons from the products of glycolysis, link reaction and Krebs cycle It takes place in mitochondrial inner membrane to release ATP with water molecule by terminal oxidation (Figure 14.5).

14.5.1 Glycolysis

(*Gr: Glykos* = Glucose, *Lysis* = Splitting) Glycolysis is a linear series of reactions in which 6-carbon glucose is split into two molecules of 3-carbon pyruvic acid. The enzymes which are required for glycolysis are present in the cytoplasm (Figure 14.6). The reactions of glycolysis were worked out in yeast cells by three scientists **Gustav Embden** (German), **Otto Meyerhoff** (German) and **J Parnas** (Polish) and so it is also called as **EMP pathway**. It is the first and common stage for both aerobic and anaerobic respiration. It is divided into two phases.

1. **Preparatory phase** or endergonic phase or hexose phase (steps 1-5).
2. **Pay off phase** or oxidative phase or exergonic phase or triose phase (steps 6-10).




Figure 14.5: Overall stages of Respiration

1. Preparatory phase

Glucose enters the glycolysis from sucrose which is the end product of photosynthesis. Glucose is phosphorylated into glucose-6-phosphate by the enzyme hexokinase, and subsequent reactions are carried out by different enzymes (Figure 14.6). At the end of this phase fructose-1, 6 - bisphosphate is cleaved into glyceraldehyde-3- phosphate and dihydroxy acetone phosphate by the enzyme aldolase. These two are isomers. Dihydroxy acetone phosphate is isomerised into glyceraldehyde-3- phosphate by the enzyme triose phosphate isomerase, now two molecules of glyceraldehyde 3 phosphate enter into pay off phase. During preparatory phase two ATP molecules are consumed in step-1 and step-3 (Figure 14.6).

Check your grasp!
How many ATP molecules are produced from one sucrose molecule?

2. Pay off phase

Two molecules of glyceraldehyde-3-phosphate oxidatively phosphorylated into two molecules of 1,3 - bisphospho glycerate. During this reaction $2NAD^+$ is reduced to $2NADH + H^+$ by glyceraldehyde- 3- phosphate dehydrogenase at step 6. Further reactions are carried out by different enzymes and at the end two molecules of pyruvate are produced. In this phase, 2ATPs are produced at step 7 and 2 ATPs at step10 (Figure 14.6). Direct transfer of phosphate moiety from substrate molecule to ADP and is converted into ATP is called **substrate phosphorylation**

XIth_Botany_Chapter_14_EM.indd 294 5/17/2020 9:20:58 PM