2. Cortex is the region found between (NEET II – 2016)
   a. epidermis and stele
   b. pericycle and endodermis
   c. endodermis and pith
   d. endodermis and vascular bundle

3. Read I – IV and find the correct order of components from outer side to inner side in a woody dicot stem (CBSE -AIPMT – 2015)
   (I) secondary Cortex (II) wood
   (III) secondary phloem (IV) phellem
   a. III, IV, II and I b. I, II, IV and III
   c. IV, I, III and II d. IV, III, I and II

4. You are given a fairly old piece of a dicot stem and a dicot root. Which of the following anatomical structures will you use to distinguish between the two? (CBSE -AIPMT 2014)
   a. secondary xylem
   b. secondary phloem
   c. protoxylem
   d. cortical cells

5. Heart wood differs from sapwood in (CBSE -AIPMT 2010)
   a. the presence of rays and fibres
   b. the absence of vessels and parenchyma
   c. having dead and non-conducting elements
   d. being susceptible to hosts and pathogens

6. The annular and spirally thickened conducting elements generally develop in the protoxylem when the root or stem is (CBSE -AIPMT 2009)
   a. maturing b. elongating
   c. widening d. differentiating

7. Anatomically fairly old dicotyledonous root is distinguished from the dicotyledonous stem by the (CBSE- AIPMT 2009)
   a. absence of secondary xylem
   b. absence of secondary phloem
   c. presence of cortex
   d. position of protoxylem

8. In barley stem, vascular bundles are (CBSE -AIPMT 2009)
   a. open and scattered
   b. closed and scattered
   c. open and in a ring
   d. closed and radial

9. Palisade parenchyma is absent in the leaves of (CBSE- AIPMT 2009)
   a. sorghum b. mustard
   c. soyabean d. gram

10. Sugarcane plant has (AIIMS 2009)
    a. reticulate venation
    b. capsular fruits
    c. pentamerous flowers
    d. dump-bell shaped guard cells

11. Vascular tissues in flowering plants develop from (CBSE- AIPMT 2008 & JIPMER 2012)
    a. phellogen **b. plerome**
    c. periblem d. dermatogen

12. The length of different internodes in a culm of sugarcane is variable because of (CBSE -AIPMT 2008)
    a. short apical meristem
    b. position of axillary buds
    c. size of leaf lamina at the node below each internode
    d. intercalary meristems

13. Passage cells are thin-walled cells found in (CBSE -AIPMT 2007)
    a. endodermis of roots facilitating rapid transport of water from cortex to pericycle
    b. phloem elements that serve as entry points for substances for transport to other plant parts
    c. testa of seeds to enable emergence of growing embryonic axis during seed germination
    d. central region of style through which the pollen tube grows towards the ovary

14. Which one of the following is not a lateral meristem (CBSE -AIPMT 2010)
    a. interfascicular cambium
    b. phellogen
    c. intercalary meristem
    d. intrafascicular cambium

15. A common feature of vessel elements and sieve tube elements is (CBSE- AIPMT 2007)
    a. enucleate condition
    b. presence of P. Protein
    c. thick secondary wall
    d. pores on lateral walls

16. In a longitudinal section of a root, starting from the tip upward, the four zones occur in the following order (CBSE -AIPMT 2004)
    a. root cap, cell division, cell enlargement, cell maturation
    b. root cap, cell division, cell maturation, cell enlargement
    c. cell division, cell enlargement, cell maturation, root cap

XIth_Botany_Chapter_15_EM.indd 346 5/17/2020 9:22:04 PM