2.4 Bryophytes

Amphibians of Plant Kingdom

In the previous chapter, we noticed a wide range of thallus organization in Algae. Majority of them are aquatic. The development of heterotrichous habit, development of parenchyma tissue and dichotomous branching in some algae supports the view that colonization of plants in land occurred in the past. Bryophytes are simplest and most primitive plant groups descended from alga – like ancestors. They are simple embryophytes. Let us learn about the structure and reproduction of these primitive land plants called Bryophytes in detail.

Bryophytes are simplest land inhabiting cryptogams and are restricted to moist, shady habitats. They lack vascular tissue and hence called **'Non- vascular cryptogams'**. They are also called as '**amphibians of plant kingdom'** because they need water for completing their life cycle.





Shiv Ram Kashyap
(1882-1934)

Father of Indian Bryology. He published a book- 'Liverworts of Western Himalayas and Punjab Plains' He identified new genera like *Atchinsoniella, Sauchia, Sewardiella* and *Stephansoniella.*

2.4.1 General characteristic features

- The plant body of bryophyte is gametophyte and is not differentiated into root, stem and leaf like structure.
- Most of them are primitive land dwellers. Some of them are aquatic (*Riella, Ricciocarpus*).
- The gametophyte is conspicuous, long lived phase of the life cycle. Thalloid forms are present in liverworts and Hornworts. In Mosses leaf like, stem like structures are present. In Liverworts thallus grows prostrate on the ground and is attached to the substratum by means of rhizoids. Two types of rhizoids are present namely smooth walled and pegged or tuberculate. Multicellular scales are also present. In Moss the plant body is erect with central axis bearing leaf like expansions. Multicellular rhizoids are present. The structure and reproduction in Bryophytes is given in Figure 2.8
- Vascular tissue like xylem and phloem are completely absent, hence called 'Non vascular cryptogams'.
- Vegetative reproduction takes place by the formation of adventitious buds (*Riccia fluitans*) tubers develop in *Anthoceros.* In some forms small detachable branches or brood bodies are formed, they help in vegetative reproduction as in *Bryopteris fruticulosa*. In *Marchantia* propagative organs called gemmae are formed and help in reproduction.
- Sexual reproduction is oogamous. Antheridia and Archegonia are produced in a protective covering and are multicellular.
- The antheridia produces biflagellate antherozoids which swims in thin film of water and reach the archegonium and fuse with the egg to form diploid zygote.
- Water is essential for fertilization.
- The zygote is the first cell of the sporophyte generation. It undergoes mitotic division to form multicellular undifferentiated embryo. The embryogeny is exoscopic (the first division of the zygote is transverse and the apex of the embryo

XIth_Botany_Chapter_2_EM.indd 41 5/17/2020 9:01:05 PM