2. Trailers (Stragglers)

It is a weak stem that spreads over the surface of the ground without rooting at nodes. They are divided into 3 types,

i. **Prostrate (Procumbent):** A stem that grows flat on the ground. Example: *Indigofera prostrata.*

ii. **Decumbent:** A stem that grows flat but becomes erect during reproductive stage. Example: *Tridax.*

iii. **Diffuse:** A trailing stem with spreading branches. Example: *Boerhavia diffusa.*

3. Climbers

These plants have long weak stem and produce special organs for attachment for climbing over a support. Climbing helps to display the leaves towards sunlight and to position the flower for effective pollination.

i. Root climbers

Plants climbing with the help of adventitious roots (arising from nodes) as in species of *Piper betel, Piper nigrum, Pothos.*

ii. Stem climbers (Twiners)

These climbers lack specialised structure for climbing and the stem itself coils around the support. Example: *Ipomoea, Clitoria, Quisqualis.*

Stem climbers may coil around the support either clockwise or anti-clockwise. Clockwise coiling climbers are called **dextrose.** Example: *Dioscorea alata*. Anti-clockwise coiling climbers are called **sinistrose.** Example: *Dioscorea bulbifera.*

iii. Hook climbers

These plants produce specialized hook like structures which are the modification of various organs of the plant. In *Artabotrys* inflorescence axis is modified into hook. In *Calamus* (curved hook) leaf tip is modified into hook. In *Bignonia unguis-cati* the leaflets are modified into curved hook (figure: 3.17). In *Hugonia* the axillary buds modified into hook.

iv. Thorn climbers

Climbing or reclining on the support with the help of thorns as in *Bougainvillea* and *Carissa.*

v. Lianas (woody stem climber)

Woody perennial climbers found in tropical forests are lianas. They twine themselves around tall trees to get light. Example*: Hiptage benghalensis, Bauhinia vahlii.*

vi. Tendril climbers

Tendrils are thread-like coiling structures which help the plants in climbing. Tendrils may be modifications of Stem – as in *Vitis* and *Cissus quadrangularis;* Inflorescence axis – *Antigonon;* Leaf – *Lathyrus;* Leaflets - *Pisum sativum;* Petiole – *Clematis;* Leaftip – *Gloriosa;* Stipules – *Smilax*. In pitcher plant (*Nepenthes*) the midrib of the leaf often coils around a support like a tendril and holds the pitcher in a vertical position.

4. Phylloclade

This is a green, flattened cylindrical or angled stem or branch of unlimited growth, consisting of a series of nodes and internodes at long or short intervals. Phylloclade is characteristic adaptation of xerophytes where the leaves often fall off early and modified into spines or scales to reduce transpiration. The phylloclade takes over all the functions of leaves, particularly photosynthesis. The phylloclade is also called as **cladophyll.** Example: *Opuntia, Phyllocactus, Muehlenbeckia* (flattened phylloclade) *Casuarina, Euphorbia tirucalli, Euphorbia antiquorum* (cylindrical phylloclade).




Figure 3.8: (a) Phylloclade-*Opuntia* (b) Cladode-*Asparagus*

5. Cladode

Cladode is a flattened or cylindrical stem similar to Phylloclade but with one or two

XIth_Botany_Chapter_3_EM.indd 68 5/17/2020 9:02:37 PM