Vegetarians use meat replacements made of textured vegetable protein (soy Protein)

Folic acid is also called pteroylglutamic acid, a deficiency of which results in macrocytic anaemia.

6.2 NUTRITIONAL NEEDS DURING DIFFERENT STATGES OF HUMAN LIFE CYCLE

6.2.1 Nutritional Needs of Infants

Infancy is a period of rapid growth. The development during infancy is more rapid than during at any other period in the life time of an individual.

Growth and Development

The growth and development is accompanied by a number of physiological changes which include changes in body size and body composition, changes in the gastro intestinal system, excretory system and circulatory system. Nutrition is crucial and proper dietary modifications are vital during this period.

Immunization

Malnutrition during infancy leads to a higher incidence of infant mortality. Besides malnutrition,infection causes mortality. Immunization protects the children against disease. Table 6 below presents the immunization schedule for infants and children.

Good nutrition is essential for the growth and development that occurs during an infant's first year of life. As an infant's mouth, tongue, and digestive tract mature, the infant shifts from being able to only suckle, swallow, and take in liquid foods, such as breast milk or infant formula, to

Table 6 National Immunization Schedule (NIS) for Infants, Children

Age	Vaccine
Soon after birth	Hepatitis B 1st dose, OPV 1st dose, BCG
6 weeks	Hepatitis B 2nddose,DPT 1stdose,OPV 2nd dose
10 weeks	DPT 2nddose,OPV 2nd dose
14 weeks	DPT 3rddose,OPV 4th dose
6 weeks, 10 weeks and 14 weeks	OPV 5thdose,Hepatitis B 3rd dose
9 months to 12 months	Measles
15 to 18 months	MMR
16-24 months	DPT, OPV 1st Booster
2 years	Typhoid vaccine
5 – 6 years	DPT, OPV 2nd Booster

DPT- Diphtheria, Pertussis, And Tetanus: OPV-Oral Polio Vaccine; MMR- Measles, mumps, and rubella

Source: https://mohfw.gov.in/sites/default/files/24545352106148966387 3.pdf

Chapter6_120-155.indd 128 31-03-2018 13:24:57