
Table 1 (continued)

	Middle age	Forty to sixty years
	Old age	After sixty years

Activity 2

Collect photos/ Pictures of each stage of the life span and paste them on a chart. Observe and record pattern of development.

ii. Period of the Neonate – from cutting and tying of the umbilical cord to approximately the end of the second week of postnatal life. During this period, the infant must make adjustments to the new environment.

a. Physical Development

Infants differ greatly in appearance and physiological functions at birth and in their early adjustments after birth.

- **Body Length and Weight**
  At birth, the average infant weighs 3 to 3.5 kg and measures 50 cms (20 inches) in length. Weight in relation to height is less at birth. The muscles of the newborn are soft, small and uncontrolled.

- **Physical Proportions**
  Babies have a large head and prominent forehead, chubby cheeks and the chin is much too small. The nose is small and flat. The neck is short and almost invisible.
  The shoulders are narrow, while the abdomen is large and bulging. Hands and feet are miniature. With the birth cry, the lungs are inflated and respiration begins. Neonatal heart beat is more rapid. Elimination of waste products begins in a few hours after birth. The relative proportion is shown in figure 2.

b. Motor Activities

The first activities of human infant are random, imperfect and uncoordinated. The most common of these include visual fixation of light, spontaneous eye movement, sucking, swallowing, lifting the head, hand movements, leg and foot movements and body jerks.

c. Sensory Abilities of the infant

i. Sight

- At birth, the retina has not reached its full development. There is no coordination between both eyes. At about thirty hours, the papillary

Chapter 2_011-045.indd 5 18-01-2019 13:21:02