


▲ Fig. 4 Adjustments during infancy

the hospital or home may vary from 60 to 70°F. Newborns are usually kept warm by providing suitable warm clothing, warm bedroom or placed next to the mother's warm body.

iii. Sucking and Swallowing

Infant must now get nourishment by sucking and swallowing. These reflexes are perfectly developed at birth and are increased along with development.

iv. Elimination

The infant's organs of elimination begin to work soon after birth. After birth, defecation occurs at an average of 4 to 7 times and urination at an average of 18 times.

d. Emotional Development of Infancy

At birth, the emotions consist of only pleasure and pain. Pleasantness is expressed by relaxing of body and is elicited through patting, rocking, warmth and sucking. Unpleasantness is expressed by tensing of the body, kicking, wriggling and crying.

e. Cognitive Development

Newborns interact with their environment entirely through reflexive behaviours. They follow their instincts and involuntary reactions to get what they need: food, air, and attention. Babies begin to grow and learn about their environment through their senses.

2.2.2 Babyhood

Babyhood continues from infancy to the second year. It is the true foundation age with rapid growth and development.

a. Physical Development during Babyhood

- **Height**

  During babyhood, changes in the overall size of the child's body are more rapid than any other time after birth. The baby measures between 23 and 24 inches at four months, by the end of one year the baby measures between 28 and 30 inches and between 32 and 34 inches at two years.

- **Weight**

  During the first year, weight changes are more than height changes. At four months the babies weight will double their birth weight and triple it at 1 year. At one year, babies weigh, on the average, three times as much as they did at birth, or approximately 10 kilograms. Increase in weight during babyhood comes mainly from an increase in fat tissue.

- **Physical Proportions**

  Growth of the head slows down in babyhood while the trunk and limb growth increases. Thus, the baby gradually becomes less top-heavy and appears more slender and less chubby by the end of babyhood.

Chapter 2_011-045.indd 7 18-01-2019 13:21:03