By comparing the components on both sides, we get

$$T - m_1 g = -m_1 a$$

$$m_1 g - T = m_1 a \qquad (3.10)$$

Adding equations (3.9) and (3.10), we get

$$m_1 g - m_2 g = m_1 a + m_2 a$$

$$(m_1 - m_2) g = (m_1 + m_2) a \qquad (3.11)$$

From equation (3.11), the acceleration of both the masses is

$$a = \left(\frac{m_1 - m_2}{m_1 + m_2} \right) g \qquad (3.12)$$

If both the masses are equal (m_1=m_2), from equation (3.12)

$$a = 0$$

This shows that if the masses are equal, there is no acceleration and the system as a whole will be at rest.

To find the tension acting on the string, substitute the acceleration from the equation (3.12) into the equation (3.9).

$$T - m_2 g = m_2 \left(\frac{m_1 - m_2}{m_1 + m_2} \right) g$$

$$T = m_2 g + m_2 \left(\frac{m_1 - m_2}{m_1 + m_2} \right) g \qquad (3.13)$$

By taking $m_2 g$ common in the RHS of equation (3.13)

$$T = m_2 g \left(1 + \frac{m_1 - m_2}{m_1 + m_2} \right)$$

$$T = m_2 g \left(\frac{m_1 + m_2 + m_1 - m_2}{m_1 + m_2} \right)$$

$$T = \left(\frac{2 m_1 m_2}{m_1 + m_2} \right) g$$

Equation (3.12) gives only magnitude of acceleration.

For mass m_1, the acceleration vector is given by $\vec{a} = -\left(\frac{m_1 - m_2}{m_1 + m_2} \right) g \hat{j}$

For mass m_2, the acceleration vector is given by $\vec{a} = \left(\frac{m_1 - m_2}{m_1 + m_2} \right) g \hat{j}$

Case 2: Horizontal motion

In this case, mass m_2 is kept on a horizontal table and mass m_1 is hanging through a small pulley as shown in Figure 3.17. Assume that there is no friction on the surface.




Figure 3.17 Blocks in horizontal motion

UNIT-3(XI-Physics_Vol-1).indd 127 28-01-2020 13:48:09