body moves with the initial velocity of the first body.

Case 3:

The first body is very much lighter than the second body

$\left(m_1 \ll m_2, \frac{m_1}{m_2} \ll 1\right)$ then the ratio $\frac{m_1}{m_2} \approx 0$ and also if the target is at rest $(u_2 = 0)$

Dividing numerator and denominator of equation (4.53) by m_2, we get

$$v_1 = \left(\frac{\frac{m_1}{m_2} - 1}{\frac{m_1}{m_2} + 1}\right) u_1 + \left(\frac{2}{\frac{m_1}{m_2} + 1}\right)(0)$$

$$v_1 = \left(\frac{0-1}{0+1}\right) u_1$$

$$v_1 = -u_1 \qquad (4.59)$$

Similarly,

Dividing numerator and denominator of equation (4.54) by m_2, we get

$$v_2 = \left(\frac{2\frac{m_1}{m_2}}{\frac{m_1}{m_2} + 1}\right) u_1 + \left(\frac{1 - \frac{m_1}{m_2}}{\frac{m_1}{m_2} + 1}\right)(0)$$

$$v_2 = (0) u_1 + \left(\frac{1 - \frac{m_1}{m_2}}{\frac{m_1}{m_2} + 1}\right)(0)$$

$$v_2 = 0 \qquad (4.60)$$

The equation (4.59) implies that the first body which is lighter returns back (rebounds) in the opposite direction with the same initial velocity as it has a negative sign. The equation (4.60) implies that the second body which is heavier in mass continues to remain at rest even after collision. For example, if a ball is thrown at a fixed wall, the ball will bounce back from the wall with the same velocity with which it was thrown but in opposite direction.

Case 4:

The second body is very much lighter than the first body

$\left(m_2 \ll m_1, \frac{m_2}{m_1} \ll 1\right)$ then the ratio $\frac{m_2}{m_1} \approx 0$ and also if the target is at rest $(u_2 = 0)$

Dividing numerator and denominator of equation (4.53) by m_1, we get

$$v_1 = \left(\frac{1 - \frac{m_2}{m_1}}{1 + \frac{m_2}{m_1}}\right) u_1 + \left(\frac{2\frac{m_2}{m_1}}{1 + \frac{m_2}{m_1}}\right)(0)$$

$$v_1 = \left(\frac{1-0}{1+0}\right) u_1 + \left(\frac{0}{1+0}\right)(0)$$

$$v_1 = u_1 \qquad (4.61)$$

Similarly,

Dividing numerator and denominator of equation (4.58) by m_1, we get

$$v_2 = \left(\frac{2}{1 + \frac{m_2}{m_1}}\right) u_1 + \left(\frac{\frac{m_2}{m_1} - 1}{1 + \frac{m_2}{m_1}}\right)(0)$$

$$v_2 = \left(\frac{2}{1+0}\right) u_1$$

$$v_2 = 2u_1 \qquad (4.62)$$

UNIT-4(XI-Physics_Vol-1).indd 197 28-01-2020 13:52:07