$$= \frac{(v_2 - v_1)}{(u_1 - u_2)} = \frac{(v_2 - v_1)}{(u_1 - 0)} = \frac{(v_2 - v_1)}{u_1}$$

$$\Rightarrow v_2 - v_1 = e\,u_1 \qquad (1)$$

From the law of conservation of linear momentum,

$$m\,u_1 = m\,v_1 + m\,v_2 \Rightarrow u_1 = v_1 + v_2 \quad (2)$$

Using the equation (2) for u_1 in (1), we get

$$v_2 - v_1 = e(v_1 + v_2)$$

On simplification, we get

$$\frac{v_1}{v_2} = \frac{1-e}{1+e}$$

SUMMARY

- When a force $\vec{F}$ acting on an object displaces it by $d\vec{r}$, then the work done(W) by the force is $W = \vec{F}.d\vec{r} = Fdr\cos\theta$.
- The work done by the variable force is defined by $\int_i^f \vec{F}.d\vec{r}$
- Work-kinetic energy theorem: The work done by a force on the object is equal to the change in its kinetic energy.
- The kinetic energy can also be defined in terms of momentum which is given by $K.E = \frac{p^2}{2m}$.
- The potential energy at a point P is defined as the amount of work required to move the object from some reference point O to the point P with constant velocity. It is given by $U = \int_O^P \vec{F}_{ext}.d\vec{r}$. The reference point can be taken as zero potential energy.
- The gravitational potential energy at a height h is given by $U = mgh$. When the elongation or compression is x, the spring potential energy is given by $U = \frac{1}{2}kx^2$. Here k is spring constant.
- The work done by a conservative force around the closed path is zero and for a non-conservative force it is not zero.
- The gravitational force, spring force and Coulomb force are all conservative but frictional force is non-conservative.
- In the conservative force field, the total energy of the object is conserved.
- In the vertical circular motion, the minimum speed required by the mass to complete the circle is $\sqrt{5gr}$. Where r , is the radius of the circle.
- Power is defined as the rate of work done or energy delivered. It is equal to $P = \frac{W}{t} = \vec{F}.\vec{v}$
- The total linear momentum of the system is always conserved for both the elastic and inelastic collisions.
- The kinetic energy of the system is conserved in elastic collisions.
- The coefficient of restitution = $\frac{velocity\ of\ separation(after\ collision)}{velocity\ of\ approach(before\ collision)}$

UNIT-4(XI-Physics_Vol-1).indd 201 28-01-2020 13:52:17