


Figure 5.21 Moment of inertia of uniform rod

the rod. First an origin is to be fixed for the coordinate system so that it coincides with the centre of mass, which is also the geometric centre of the rod. The rod is now along the x axis. We take an infinitesimally small mass (dm) at a distance (x) from the origin. The moment of inertia (dI) of this mass (dm) about the axis is,

$$dI = (dm)\,x^2$$

As the mass is uniformly distributed, the mass per unit length (λ) of the rod is, $\lambda = \frac{M}{\ell}$

The (dm) mass of the infinitesimally small length as, $dm = \lambda\, dx = \frac{M}{\ell} dx$

The moment of inertia (I) of the entire rod can be found by integrating dI,

$$I = \int dI = \int (dm)\,x^2 = \int \left(\frac{M}{\ell} dx\right) x^2$$

$$I = \frac{M}{\ell}\int x^2 dx$$

As the mass is distributed on either side of the origin, the limits for integration are taken from $-\ell/2$ to $\ell/2$.

$$I = \frac{M}{\ell}\int_{-\ell/2}^{\ell/2} x^2 dx = \frac{M}{\ell}\left[\frac{x^3}{3}\right]_{-\ell/2}^{\ell/2}$$

$$I = \frac{M}{\ell}\left[\frac{\ell^3}{24} - \left(-\frac{\ell^3}{24}\right)\right] = \frac{M}{\ell}\left[\frac{\ell^3}{24} + \frac{\ell^3}{24}\right]$$

$$I = \frac{M}{\ell}\left[2\left(\frac{\ell^3}{24}\right)\right]$$


$$I = \frac{1}{12} M\ell^2 \qquad (5.41)$$

EXAMPLE 5.14

Find the moment of inertia of a uniform rod about an axis which is perpendicular to the rod and touches any one end of the rod.

Solution

The concepts to form the integrand to find the moment of inertia could be borrowed from the earlier derivation. Now, the origin is fixed to the left end of the rod and the limits are to be taken from 0 to ℓ.



$$I = \frac{M}{\ell}\int_0^{\ell} x^2 dx = \frac{M}{\ell}\left[\frac{x^3}{3}\right]_0^{\ell} = \frac{M}{\ell}\left[\frac{\ell^3}{3}\right]$$

$$I = \frac{1}{3} M\ell^2$$

UNIT-5(XI-Physics_Vol-1).indd 235 28-01-2020 13:56:07