If we take $x_1 = 2$ and $x_2 = 2.25$, then $y_1 = 4$ and $y_2 = 5.0625$

$$\text{Here } \Delta x = 0.25 = \frac{1}{4},\ \Delta y = 1.0625$$

$$\frac{\Delta y}{\Delta x} = \frac{5.0625 - 4}{0.25} = \frac{(5.0625 - 4)}{\frac{1}{4}}$$

$$= 4(5.0625 - 4) = 4.25$$

If we take $x_1 = 2$ and $x_2 = 2.1$, then $y_1 = 4$ and $y_2 = 4.41$

$$\text{Here } \Delta x = 0.1 = \frac{1}{10} \text{ and}$$

$$\frac{\Delta y}{\Delta x} = \frac{(4.41 - 4)}{\frac{1}{10}} = 10(4.41 - 4) = 4.1$$

These results are tabulated as shown below:

x_1	x_2	Δx	y_1	y_2	$\frac{\Delta y}{\Delta x}$
2	2.25	0.25	4	5.0625	4.25
2	2.1	0.1	4	4.41	4.1
2	2.01	0.01	4	4.0401	4.01
2	2.001	0.001	4	4.004001	4.001
2	2.0001	0.0001	4	4.00040001	4.0001

From the above table, the following inferences can be made.

- As Δx tends to zero, $\frac{\Delta y}{\Delta x}$ approaches the limit given by the number 4.
- At a point x = 2, the derivative $\frac{dy}{dx} = 4$.
- It should also be mentioned here that $\Delta x \to 0$ does not mean that $\Delta x = 0$.

This is because, if we substitute $\Delta x = 0$, $\frac{\Delta y}{\Delta x}$ becomes indeterminate.

In general, we can obtain the derivative of the function $y = x^2$, as follows:

$$\frac{\Delta y}{\Delta x} = \frac{(x + \Delta x)^2 - x^2}{\Delta x} = \frac{x^2 + 2x\Delta x + \Delta x^2 - x^2}{\Delta x}$$

$$= \frac{2x\Delta x + \Delta x^2}{\Delta x} = 2x + \Delta x$$

$$\frac{dy}{dx} = \lim_{\Delta x \to 0} 2x + \Delta x = 2x$$

The table below shows the derivatives of some common functions used in physics

Function	Derivative
$y = x$	$dy/dx = 1$
$y = x^2$	$dy/dx = 2x$
$y = x^3$	$dy/dx = 3x^2$
$y = x^n$	$dy/dx = nx^{n-1}$
$y = \sin x$	$dy/dx = \cos x$
$y = \cos x$	$dy/dx = -\sin x$
y = constant	$dy/dx = 0$
$y = AB$	$\frac{dy}{dx} = A\left(\frac{dB}{dx}\right) + \left(\frac{dA}{dx}\right)B$

In physics, velocity, speed and acceleration are all derivatives with respect to time 't'. This will be dealt with in the next section.

EXAMPLE 2.19

Find the derivative with respect to t, of the function $x = A_0 + A_1 t + A_2 t^2$ where A_0, A_1 and A_2 are constants.

Solution

Note that here the independent variable is 't' and the dependent variable is 'x'

UNIT-2(XI-Physics_Vol-1).indd 63 28-01-2020 13:45:28