the structure and molecular nature of the system. Unlike solids and liquids, gases have two specific heats: specific heat capacity at constant pressure (s_p) and specific heat capacity at constant volume (s_v).

8.7.1 Specific heat capacity

Specific heat capacity at constant pressure (s_p):

The amount of heat energy required to raise the temperature of one kg of a substance by 1 K or 1°C by keeping the pressure constant is called specific heat capacity of at constant pressure. When the heat energy is supplied to the gas, it expands to keep the pressure constant as shown in Figure 8.23




Figure 8.23 Specific heat capacity at constant pressure

In this process a part of the heat energy is used for doing work (expansion) and the remaining part is used to increase the internal energy of the gas.

Specific heat capacity at constant volume (s_v):

The amount of heat energy required to raise the temperature of one kg of a substance by 1 K or 1°C by keeping the volume constant is called specific heat capacity at constant volume.

If the volume is kept constant, then the supplied heat is used to increase only the internal energy. No work is done by the gas as shown in Figure 8.24.




Figure 8.24 Specific heat capacity at constant volume

It implies that to increase the temperature of the gas at constant volume requires less heat than increasing the temperature of the gas at constant pressure. In other words s_p is always greater than s_v.

Molar Specific heat capacities

Sometimes it is useful to calculate the molar heat capacities C_p and C_v. The amount of heat required to raise the temperature of one mole of a substance by 1K or 1°C at constant volume is called molar specific heat capacity at constant volume (C_v). If pressure is kept constant, it is called molar specific heat capacity at constant pressure (C_p).

UNIT-8(XI-Physics_Vol-2).indd 123 17-08-2020 16:17:18