$$F_1 \propto l \Rightarrow F_1 = -kl \qquad (10.28)$$

Substituting equation (10.28) in equation (10.27), we get

$$-kl + mg = 0$$
$$mg = kl$$
or
$$\frac{m}{k} = \frac{l}{g} \qquad (10.29)$$

Suppose we apply a very small external force on the mass such that the mass further displaces downward by a displacement y, then it will oscillate up and down. Now, the restoring force due to this stretching of spring (total extension of spring is $y + l$) is

$$F_2 \propto (y + l)$$
$$F_2 = -k(y + l) = -ky - kl \qquad (10.30)$$

Since, the mass moves up and down with acceleration $\frac{d^2y}{dt^2}$, by drawing the free body diagram for this case, we get

$$-ky - kl + mg = m\frac{d^2y}{dt^2} \qquad (10.31)$$

The net force acting on the mass due to this stretching is

$$F = F_2 + mg$$
$$F = -ky - kl + mg \qquad (10.32)$$

The gravitational force opposes the restoring force. Substituting equation (10.29) in equation (10.32), we get

$$F = -ky - kl + kl = -ky$$

Applying Newton's law, we get

$$m\frac{d^2y}{dt^2} = -ky$$
$$\frac{d^2y}{dt^2} = -\frac{k}{m}y \qquad (10.33)$$

The above equation is in the form of simple harmonic differential equation. Therefore, we get the time period as

$$T = 2\pi\sqrt{\frac{m}{k}} \text{ second} \qquad (10.34)$$

Note

The time period obtained for horizontal oscillations of spring and for vertical oscillations of spring are found to be equal.

The time period can be rewritten using equation (10.29)

$$T = 2\pi\sqrt{\frac{m}{k}} = 2\pi\sqrt{\frac{l}{g}} \text{ second} \qquad (10.35)$$

The acceleration due to gravity g can be computed from the formula

$$g = 4\pi^2\left(\frac{l}{T^2}\right) \text{ m s}^{-2} \qquad (10.36)$$

EXAMPLE 10.8

A spring balance has a scale which ranges from 0 to 25 kg and the length of the scale is 0.25m. It is taken to an unknown planet X where the acceleration due to gravity is 11.5 m s^{-1}. Suppose a body of mass M kg is suspended in this spring and made to oscillate with a period of 0.50 s. Compute the gravitational force acting on the body.

Solution

Let us first calculate the stiffness constant of the spring balance by using equation (10.29),

$$k = \frac{mg}{l} = \frac{25 \times 11.5}{0.25} = 1150\ N\,m^{-1}$$

The time period of oscillations is given by $T = 2\pi\sqrt{\frac{M}{k}}$, where M is the mass of the body.

Since, M is unknown, rearranging, we get

UNIT-10(XI-Physics_Vol-2).indd 202 17-08-2020 16:23:43