remains the same during the propagation. After some time t, the pulse moving towards the right and any point on it can be represented by x' (read it as x prime) as shown in Figure 11.23 (b). Then,

$$y(x, t) = f(x') = f(x - vt) \qquad (11.35)$$

Similarly, if the wave pulse moves towards left with constant speed v, then $y = f(x + vt)$. Both waves $y = f(x + vt)$ and $y = f(x - vt)$ will satisfy the following one dimensional differential equation known as the wave equation

$$\frac{\partial^2 y}{\partial x^2} = \frac{1}{v^2}\frac{\partial^2 y}{\partial t^2} \qquad (11.36)$$

where the symbol ∂ represents partial derivative (read $\frac{\partial y}{\partial x}$ as partial y by partial x). Not all the solutions satisfying this differential equation can represent waves, because any physical acceptable wave must take finite values for all values of x and t. But if the function represents a wave then it must satisfy the differential equation. Since, in one dimension (one independent variable), the partial derivative with respect to x is the same as total derivative in coordinate x, we write

$$\frac{d^2 y}{dx^2} = \frac{1}{v^2}\frac{d^2 y}{dt^2} \qquad (11.37)$$


This can be extended to more than one dimension (two, three, etc.). Here, for simplicity, we focus only on the one dimensional wave equation.

EXAMPLE 11.11

Sketch $y = x - a$ for different values of a.

Solution


This implies, when increasing the value of a, the line shifts towards right side. For $a = vt$, $y = x - vt$ satisfies the differential equation. Though this function satisfies the differential equation, it is not finite for all values of x and t. Hence, it does not represent a wave.



EXAMPLE 11.12

How does the wave $y = \sin(x - a)$ for $a = 0$, $a = \frac{\pi}{4}$, $a = \frac{\pi}{2}$, $a = \frac{3\pi}{2}$ and $a = \pi$ look like?. Sketch this wave.

Solution



From the above picture we observe that $y = \sin(x - a)$ for $a = 0$, $a = \frac{\pi}{4}$, $a = \frac{\pi}{2}$, $a = \frac{3\pi}{2}$ and $a = \pi$, the function $y = \sin(x - a)$ shifts towards right. Further, we can take $a = vt$ and $v = \frac{\pi}{4}$, and sketching for different

UNIT-11(XI-Physics_Vol-2)_14-08-2020.indd 244 17-08-2020 15:30:57