EXAMPLE 6.7

1. Calculate the value of g in the following two cases:
   (a) If a mango of mass ½ kg falls from a tree from a height of 15 meters, what is the acceleration due to gravity when it begins to fall?

Solution

$$g' = g\left(1 - 2\frac{h}{R_e}\right)$$

$$g' = 9.8\left(1 - \frac{2\times 15}{6400\times 10^3}\right)$$

$$g' = 9.8\left(1 - 0.469\times 10^{-5}\right)$$

But $1 - 0.00000469 \cong 1$

Therefore $g' = g$

(b) Consider a satellite orbiting the Earth in a circular orbit of radius 1600 km above the surface of the Earth. What is the acceleration experienced by the satellite due to Earth's gravitational force?

Solution

$$g' = g\left(1 - 2\frac{h}{R_e}\right)$$

$$g' = g\left(1 - \frac{2\times 1600\times 10^3}{6400\times 10^3}\right)$$

$$g' = g\left(1 - \frac{2}{4}\right)$$

$$g' = g\left(1 - \frac{1}{2}\right) = g/2$$

The above two examples show that the acceleration due to gravity is a constant near the surface of the Earth.

Note

Can we substitute h = R_e in the equation 6.46? No. To get equation 6.46 we assumed that h << R_e. However h = R_e can be substituted in equation 6.45.

Variation of g with depth:

Consider a particle of mass m which is in a deep mine on the Earth. (Example: coal mines in Neyveli). Assume the depth of the mine as d. To calculate g' at a depth d, consider the following points.




Figure 6.17(b) Particle in a mine

The part of the Earth which is above the radius $(R_e - d)$ do not contribute to the acceleration. The result is proved earlier and is given as

$$g' = \frac{GM'}{(R_e - d)^2} \qquad (6.47)$$

Here M' is the mass of the Earth of radius $(R_e - d)$

Assuming the density of Earth ρ to be constant,

$$\rho = \frac{M}{V} \qquad (6.48)$$

UNIT-6(XI-Physics_Vol-2).indd 20 17-08-2020 16:15:32