Therefore,

$$g' = g - \omega^2 R \cos^2 \lambda \qquad (6.52)$$

From the expression (6.52), we can infer that at equator, $\lambda = 0$; $g' = g - \omega^2 R$. The acceleration due to gravity is minimum. At poles $\lambda = 90°$; $g' = g$, it is maximum. At the equator, g' is minimum.

EXAMPLE 6.8

Find out the value of g' in your school laboratory?

Solution

Calculate the latitude of the city or village where the school is located. The information is available in Google search. For example, the latitude of Chennai is approximately 13 degree.

$$g' = g - \omega^2 R \cos^2 \lambda$$

Here $\omega^2 R = (2\text{x}3.14/86400)^2 \text{ x } (6400\text{x}10^3) = 3.4\text{x}10^{-2}$ m s^{-2}.

It is to be noted that the value of λ should be in radian and not in degree. 13 degree is equivalent to 0.2268 rad.

$$g' = 9.8 - \left(3.4 \times 10^{-2}\right) \times \left(\cos 0.2268\right)^2$$

$$g' = 9.7677 \text{ m s}^{-2}$$

Points to Contemplate

Suppose you move towards east-west along the same latitude. Will the value of g' change?

6.4 ESCAPE SPEED AND ORBITAL SPEED

Hydrogen and helium are the most abundant elements in the universe but Earth's atmosphere consists mainly of nitrogen and oxygen. The following discussion brings forth the reason why hydrogen and helium are not found in abundance on the Earth's atmosphere. When an object is thrown up with some initial speed it will reach a certain height after which it will fall back to Earth. If the same object is thrown again with a higher speed, it reaches a greater height than the previous one and falls back to Earth. This leads to the question of what should be the speed of an object thrown vertically up such that it escapes the Earth's gravity and would never come back.

Consider an object of mass M on the surface of the Earth. When it is thrown up with an initial speed v_i, the initial total energy of the object is

$$E_i = \frac{1}{2} M v_i^2 - \frac{GMM_E}{R_E} \qquad (6.53)$$

where, M_E is the mass of the Earth and R_E- the radius of the Earth. The term $-\frac{GMM_E}{R_E}$ is the potential energy of the mass M.

When the object reaches a height far away from Earth and hence treated as approaching infinity, the gravitational potential energy becomes zero $[U(\infty) = 0]$ and the kinetic energy becomes zero as well. Therefore the final total energy of the object becomes zero. This is for minimum energy and for minimum speed to escape. Otherwise Kinetic energy can be nonzero.

UNIT-6(XI-Physics_Vol-2).indd 22 17-08-2020 16:15:38