Try these

Convert the following statements into linear equations:

1. On subtracting 8 from the product of 5 and a number, I get 32.
2. The sum of three consecutive integers is 78.
3. Peter had a Two hundred rupee note. After buying 7 copies of a book he was left with ₹60.
4. The base angles of an isosceles triangle are equal and the vertex angle measures 80°.
5. In a triangle ABC, ∠A is 10° more than ∠B. Also ∠C is three times ∠A. Express the equation in terms of angle B.

3.8.4 Solution of a linear equation


The value which replaces a variable in an equation so as to make the two sides of the equation equal is called a solution or root of the equation.

Example : $2x = 10$

We find that the equation is "satisfied" with the value $x = 5$. That is, if we put $x = 5$, in the equation, the value of the LHS will be equal to the RHS. Thus $x = 5$ is **a solution** of the equation. Note that no other value for x satisfies the equation. Thus one can say $x = 5$ is "**the**" solution of the equation.


(i) The DO-UNDO Method:

Statement (given)	You  Think
5 less than twice a number is 11.	The number needed is unknown. Let it be x. Twice the number gives $2x$. 5 less is $2x - 5$. This result is given to be 11.



This formation of equation can be visualized as follows:

From the number x, we reached $2x - 5$ by performing operations like subtraction, multiplication etc. So when $2x - 5 = 11$ is given, to get back to the value of x, we have to *'undo'* all that we did! Thus, we *'do'* to form the equation and *'undo'* to get the solution.



03_8th_Maths_Algebra_Chap 3.indd 100 2/13/2020 6:10:09 PM