Activity

1. Collect the blood group of your classmates. Complete the table and analyse.

Blood group	Tally marks	No. of students
A+		
B+		
AB+		
O+		
A-		
B-		
AB-		
O-		

2. Observe the last alphabet in the name of your classmates, tabulate them and answer the following questions.

Alphabet	Tally marks	No. of students (f)

1. In which letter do the names end the most?
2. In which letter do the names end the least?
3. What are the letters in which the names do not end with?
4. Girl names mostly end with _____ letter(s).
5. Boy names mostly end with _____ letter(s).

6.2.2 Construction of frequency distribution table for grouped data

Grouped data or Continuous Data:

A grouped data is any value within a certain interval. The data can take values between certain range with the highest and the lowest value. Continuous data can be tabulated in what is called as frequency distribution. They can be graphically represented using Histograms.

Example: 1. The age of persons in a village.
2. The height and the weight of the students of your class.

Now, we will consider a situation, if we collect data of marks for 50 students, it becomes very difficult to put tally for each and every marks of all the 50 students. Because if we arrange the marks in a table, it will be very large in length and not understandable at once. In this case, we use class intervals. In this table, consider the groups of data in the form of class intervals to tally the frequency for the given data.

Class Interval:

The range of the variable is grouped into number of classes, and each group is known as class interval (C.I). The difference between the upper limit (U) and the lower limit (L) of the class is known as class size.

i.e. C.I = Upper limit – Lower limit

06_8th_Maths_Statistics_Chap 6.indd 214 2/12/2020 12:20:31 PM