### 6.4.1 (i) Construction of a histogram for continuous frequency distribution:

**Example 6.6**


Draw a histogram for the following table which represents the age groups from 100 people in a village.

| **Ages** | 0-10 | 10-20 | 20-30 | 30-40 | 40-50 | 50-60 | 60-70 | 70-80 | 80-90 |
|---|---|---|---|---|---|---|---|---|---|
| **Number of people** | 11 | 9 | 8 | 20 | 25 | 10 | 8 | 6 | 3 |

***Solution:***

The given data is a continuous frequency distribution. The class intervals are drawn on x-axis and their respective frequencies on y-axis. Classes (ages) and its frequencies (number of people) are taken together to form a rectangle.

The histogram is constructed as given below.




**Note**

If class intervals do not start from '0' then, it is indicated by drawing a kink (Zig-Zag) mark (⩘) on the x-axis near the origin. If necessary, the kink mark (⩘) may be made on y-axis or on both the axes. i.e it indicates that we do not have data starting from the origin (O)

### 6.4.1 (ii) Construction of histogram for discontinuous frequency distribution:

**Example 6.7**

The following table gives the number of literate females in the age group 10 to 45 years in a town.

| **Age group** | 10-15 | 16-21 | 22-27 | 28-33 | 34-39 | 40-45 |
|---|---|---|---|---|---|---|
| **No. of females** | 350 | 920 | 850 | 480 | 230 | 200 |

Draw a histogram to represent the above data



06_8th_Maths_Statistics_Chap 6.indd 223 2/12/2020 12:20:40 PM