4. MEASUREMENT OF VOLUME OF LIQUIDS

Aim:

To measure the volume of given colourless and coloured liquids.

QLH74T

Materials required:

Pipette (20ml), sample liquids and beakers

Procedure:

Take a 20 ml pipette. Wash it thoroughly with water and then rinse it with the given liquid. Insert the lower end of the pipette into the given liquid and suck the solution slowly till the solution rises well above the circular mark on the stem. Take the pipette out of the mouth and quickly close it with the fore finger. Take the pipette out the liquid and keep it such a way that the circular mark on the stem is at the level of the eyes. Now slowly release the fore finger to let the liquid drop out until the lower meniscus touches the circular mark on the stem. Now the liquid in the pipette is exactly 20 ml. This can be transferred to an empty beaker by removing the fore finger.

Tabulation

Sl. No.	Name of the liquid	Colour of the liquid	Nature of the meniscus	Volume of the liquid
1				
2				
3				

Report: Exactly 20 ml of various liquids are measured using a standard 20 ml pipette.

Note:

1. Keeping the circular mark on the stem of the pipette above or below the level of the eyes will lead to error.
2. When colored liquids are measured, the upper meniscus should be taken into account.
3. Never suck strong acids or strong alkalis using a pipette.

5. IDENTIFICATION OF ADAPTATIONS IN ANIMALS

Aim:

To identify the given vertebrate animals and list out the adaptations seen in them.

QM2FHJ

Required specimens:

1. Pisces (Fish), 2. Amphibian (Frog), 3. Reptile (Calotes), 4. Aves (Dove), 5. Mammal (Rat)

IX_SCI_EM_Practical.indd 314 14-02-2020 11:10:46