Cushing's syndrome is caused due to excess secretion of cortisol. Obesity of the face and trunk, redness of face, hand, feet, thin skin, excessive hair growth, loss of minerals from bone (osteoporosis) systolic hypertension are features of Cushing's syndrome. Suppression of sexual function like atrophy of gonads are the other symptoms of Cushing's syndrome. (Figure 11.16)




Figure.11.16 Cushing's syndrome

Hypoglycaemia is due to increased secretion of insulin thereby blood glucose level decreases. In this disorder blood glucose level lowers than normal fasting index. Increased heartbeat, weakness, nervousness, headache, confusion, lack of co-ordination, slurred speech, serious brain defects like epilepsy and coma occurs.

Normal blood glucose level:

Preprandial : 70 – 110 mg/dl (Before food) – (Fasting)

Postprandial : 110 – 140 mg/dl (About two hours after food)

Hyperglycaemia is otherwise known as **Diabetes mellitus**. It is caused due to reduced secretion of insulin. As the result, blood glucose level is elevated. Diabetes mellitus is of two types, **Type I Diabetes** and **Type II Diabetes**. Type I diabetes is also known Insulin dependent diabetes, caused by the lack of insulin secretion due to illness or viral infections. Type II diabetes is also known as Non- Insulin dependent diabetes, caused due to reduced sensitivity to insulin, often called as insulin resistance. Symptoms of diabetes includes, polyurea (excessive urination), polyphagia (excessive intake of food), polydipsia (excessive consumption liquids due to thirst), ketosis (breakdown of fat into glucose results in accumulation of ketone bodies) in blood. Gluconeogenesis (Conversion of non- carbohydrate form like amino acids and fat into glucose) also occur in diabetes.



Avoid use of synthetic soft drinks

The branded soft drinks damage our endocrine system. While consuming soft drinks, the sugar level increases in blood which leads to elevated insulin secretion to reduce the blood glucose level. The elevated insulin level diminishes immunity and cause obesity, cardio-vascular disorders etc.

Diabates insipidus is caused due to hyposecretion of vasopressin (ADH) from neurohypophysis. The symptom includes frequent urination (polyurea) and excessive consumption of liquids due to thirst (polydipsia).

TN_GOVT_XI_Zoology_Ch11.indd 216 09-01-2020 5.20.00 PM