1.12.3 Application of Newton's law of gravitation

1) Dimensions of the heavenly bodies can be measured using the gravitation law. Mass of the Earth, radius of the Earth, acceleration due to gravity, etc. can be calculated with a higher accuracy.
2) Helps in discovering new stars and planets.
3) One of the irregularities in the motion of stars is called 'Wobble' lead to the disturbance in the motion of a planet nearby. In this condition the mass of the star can be calculated using the law of gravitation.
4) Helps to explain germination of roots is due to the property of geotropism which is the property of a root responding to the gravity.
5) Helps to predict the path of the astronomical bodies.

Points to Remember

- Mechanics is divided into statics and dynamics.
- Ability of a body to maintain its state of rest or motion is called Inertia.
- Moment of the couple is measured by the product of any one of the forces and the perpendicular distance between two forces.
- SI unit of force is newton (N). C.G.S unit is dyne.
- When a force F acts on a body for a period of time t, then the product of force and time is known as 'impulse'.
- The unit of weight is newton or kg f
- The weight of a body is more at the poles than at the equatorial region.
- Mass of a body is defined as the quantity of matter contained in the object. Its SI unit is kilogram (kg).
- Apparent weight is the weight of the body acquired due to the action of gravity and other external forces on the body.
- Whenever a body or a person falls freely under the action of Earth's gravitational force alone, it appears to have zero weight. This state is referred to as 'weightlessness'.

SOLVED PROBLEMS

Problem-1: Calculate the velocity of a moving body of mass 5 kg whose linear momentum is 2.5 $kg\,m\,s^{-1}$.

Solution: Linear momentum = mass × velocity

Velocity = linear momentum / mass.

V = 2.5 / 5 = 0.5 ms^{-1}

Problem 2: A door is pushed, at a point whose distance from the hinges is 90 cm, with a force of 40 N. Calculate the moment of the force about the hinges.

Solution:

Formula: The moment of a force M = F × d

Given: F = 40 N and d = 90 cm = 0.9 m.

Hence, moment of the force = 40 × 0.9 = 36 N m.

Problem 3 : At what height from the centre of the Earth the acceleration due to gravity will be $\frac{1}{4}^{th}$ of its value as at the Earth.

Solution:

Data: Height from the centre of the Earth, R′ = R + h

The acceleration due to gravity at that height, g′ = g/4

Formula: $g = GM/R^2$, $g' = GM/R'^2$

$$\frac{g}{g'} = \left(\frac{R'}{R}\right)^2 = \left(\frac{R+h}{R}\right)^2 = \left(1 + \frac{h}{R}\right)^2$$

$$4 = \left(1 + \frac{h}{R}\right)^2,$$

$$2 = 1 + \frac{h}{R} \qquad \text{or } h = R. \qquad R' = 2R$$

From the centre of the Earth, the object is placed at twice the radius of the earth.

10th_Science_Unit-1.indd 12 11-01-2020 11:52:13