


Figure 6.1 The Human respiratory system

gullet during normal swallowing. During swallowing a thin elastic flap called epiglottis prevents the food from entering into the larynx and avoids choking of food.

The trachea is semiflexible tube supported by multiple cartilaginous rings which extends up to the midthoracic cavity and at the level of the 5th thoracic vertebra where it divides into right and left primary bronchi, one bronchus to each lung. Within the lungs the bronchi divides repeatedly into secondary and tertiary bronchi and further divides into terminal bronchioles and respiratory bronchioles.

Bronchi have 'C' shaped curved cartilage plates to ensure that the air passage does not collapse or burst as the air pressure changes during breathing. The bronchioles are without cartilaginous rings and have rigidity that prevent them from collapsing but are surrounded by smooth muscle which contracts or relaxes to adjust the diameter of these airways.

The fine respiratory bronchioles terminate into highly vascularised thin walled pouch like air sacs called alveoli meant for gaseous exchange (Figure 6.2, 6.3). The diffusion membrane of alveolus is made up of three layers – the thin squamous epithelial cells of the alveoli, the endothelium of the alveolar capillaries and the basement substance found in between them. The thin squamous epithelial cells of the **alveoli** are composed of Type I and Type II cells. Type I cells are very thin so that gases can diffuse rapidly through them. Type II cells are thicker, synthesize and secrete a substance called **Surfactant**.

The lungs are light spongy tissues enclosed in the thoracic cavity surrounded by an airtight space. The thoracic cavity is bound dorsally by the vertebral column and ventrally by the sternum, laterally by the ribs and on the lower side by the dome shaped diaphragm.

The lungs are covered by double walled pleural membrane containing a several layers of elastic connective tissues and capillaries, which encloses the pleural fluid. Pleural fluid reduces friction when the lungs expand and contract.

Characteristic features of respiratory surface:

- surface area must be very large and richly supplied with blood vessels
- should be extremely thin and kept moist
- should be in direct contact with the environment
- should be permeable to respiratory gases

TN_Govt_Bio Zoology_XI_Std_Ch06.indd 94 09-01-2020 5.28.49 PM