Evaluation

4H9QA8

1 Which structure in the ear converts pressure waves to action potentials?

a. Tympanic membrane

b. Organ of Corti

c. Oval window

d. Semicircular canal

2. Which of the following pairings is correct?

a. Sensory nerve – afferent

b. Motor nerve - afferent

c. Sensory nerve – ventral

d. Motor nerve – dorsal

3. During synaptic transmission of nerve impulse, neurotransmitter (P) is released from synaptic vesicles by the action of ions (Q). Choose the correct P and Q.


a. P = Acetylcholine, Q = Ca++

b. P = Acetylcholine, Q = Na+

c. P = GABA, Q=Na+

d. P = Cholinesterase, Q = Ca++

4. Examine the diagram of the two cell types A and B given below and select the correct option.



a. Cell-A is the rod cell found evenly all over retina

b. Cell-A is the cone cell more concentrated in the fovea centralis

c. Cell-B is concerned with colour vision in bright light

d. Cell-A is sensitive to bright light intensities

5. Assertion: The imbalance in concentration of Na^+, K^+ and proteins generates action potential.

Reason: To maintain the unequal distribution of Na^+ and K^+, the neurons use electrical energy.

a. Both Assertion and Reason are true and Reason is the correct explanation of the Assertion.

b. Both Assertion and Reason are true but the Reason is not the correct explanations of Assertion.

c. Assertion is true, but Reason is false.

d. Both Assertion and Reason are false.

6. Which part of the human brain is concerned with the regulation of body temperature?

a. Cerebellum

b. Cerebrum

c. Medulla oblongata

d. Hypothalamus

7. The respiratory centre is present in the

a. Medulla oblongata

b. Hypothalamus

c. Cerebellum

d. Thalamus

8. Match the following human spinal nerves in column I with their respective number in column II and choose the correct option

column I	column II
P. Cervical nerves	i. 5 pairs
Q. Thoracic nerve	ii. 1 pair
R. Lumbar nerve	iii. 12 pair
S. Coccygeal nerve	iv. 8 pair

a. (P-iv),(Q-iii),(R-i),(S-ii)

b. (P-iii), (Q-i), (R-ii), (S-iv)

c. (P-iv),(Q-i),(R-ii),(S-iii)

d. (P-ii), (Q-iv), (R-i), (S-iii)

9. The abundant intracellular cation is

a. H^+ b. K^+ c. Na^+ d. Ca^{++}

TN_GOVT_XI_Bio Zoology_Ch10.indd 192 09-01-2020 5.32.00 PM