

Leghorn-Hen and Cock




White plymouth rock Hen and Cock



Silkie



Light brahma Hen and cock



Aseel - Hen and cock

Figure 12.13 Different types of chicken breeds

4. **Game breeds:** Since ancient times, special breed of roosters have been used for the sport of cockfighting.

Aseel: This breed is white or black in colour. The hens are not good egg layers but are good in incubation of eggs. It is found in all states of India. Aseel is noted for its pugnacity, high stamina, and majestic gait and dogged fighting qualities. Although poor in productivity, this breed is well-known for their meat qualities.

5. **Ornamental breeds:** Ornamental chicken are reared as pets in addition to their use for egg production and meat.

Silkie: It is a breed of chicken has a typical fluffy plumage, which is said to feel like silk and satin. The breed has numerous additional special characters, such as black skin and bones, blue earlobes, and five toes on each foot, while the majority chickens only have four. They are exhibited in poultry shows, and come out in various colours. Silkies are well recognized for their calm, friendly temperament. Silkie chicken is especially simple to maintain as pets.

Types of Poultry farming: There are different methods used to rear both broiler and layer chicken. The types of poultry farming are Free range farming, Organic method, Yarding method, Battery cage method and Furnished cage method.

Among these, Battery cage method is widely used in large scale poultry farms. The Free range, Organic and Yarding methods are eco-friendly and the eggs produced by such farming practices are preferred in the market.

Stages involved in rearing:

There are some steps involved in rearing of chicken.

1. Selection of the best layer: An active intelligent looking bird, with a bright comb, not obese should be selected.
2. Selection of eggs for hatching: Eggs should be selected very carefully. Eggs should be fertile, medium sized, dark brown shelled and freshly laid eggs are preferred for rearing. Eggs should be washed, cleaned and dried.
3. Incubation and hatching: The maintenance of newly laid eggs in optimum condition till hatching is called incubation. The fully developed chick emerges out of egg after an incubation period of 21 – 22 days. There are two types of incubation namely natural incubation and artificial incubation.In the natural incubation method, only a limited number of eggs can be incubated by a mother hen. In artificial incubation, more number of eggs can be incubated in a chamber (**Incubator**).

TN_GOVT_XI_Bio Zoology_Ch12.indd 237 09-01-2020 5.33.21 PM