
skin is smooth or rough, moist, pigmented and glandular. Eyes have eyelids and the tympanum represents the ear. Respiration is by gills, lungs and through the skin. Heart is three chambered. Kidneys are **mesonephric.** Sexes are separate and fertilization is external. They are oviparous and development is indirect. They show **hibernation** and **aestivation**.

Examples: *Bufo* (Toad), *Rana* (Frog), *Hyla* (Tree frog), *Salamandra* (Salamander), *lcthyophis* (Limbless amphibians) (Figure 2.25).



Frog *Toad*

Salamander *Icthyophis (Caecilians)*


Figure 2.25 Examples of Amphibia

2.4.8 Class: Reptilia

(*L. repere* or *reptum* – to creep or crawl)

They are mostly terrestrial animals and their body is covered by dry, and cornified skin with epidermal scales or scutes. Reptiles have three chambered heart but four chambered in crocodiles. All are cold blooded amniotes (poikilotherms). Most reptiles lay **cleidoic eggs** with extraembryonic membranes like **amnion, allantois, chorion and yolk sac**. Excretion by metanephric kidneys and are uricotelic. Sexes are separate with well marked sexual dimorphism. Internal fertilization takes place and all are oviparous.

Examples : *Chelone* (Turtle), *Testudo* (tortoise), *Hemidactylus* (House lizard), *Chameleon* (Tree lizard), *Calotes* (Garden lizard), *Draco* (Flying lizard), *Crocodilus* (crocodile), Poisonous snakes - *Naja* (Cobra), *Bangarus* (Krait), *Vipera* (Viper) (Figure 2.26).



King cobra *Draco (Flying Lizard)*

Crocodile *Chameleon*

Figure 2.26 Examples of Reptiles

TURTLE Vs TORTOISE



Turtles spend most of their life in the water
Carapace is laterally compressed and streamlined.
Mostly live in the water or are always found near it.
Most of them have webbed feet.



Tortoises spend most of their life on land.
Carapace is usually dome-shaped.
These are primarily terrestrial.
Feet are short and sturdy with bent legs.

TN_GOVT_XI_Bio Zoology_Ch02.indd 30 09-01-2020 5.24.59 PM