


Figure 5.2 Buccal cavity

involved in the process of absorption and the cells lining the villi produce numerous microscopic projections called microvilli giving a brush border appearance that increase the surface area enormously. Along with villi, the ileal mucosa also contain mucus secreting goblet cells and lymphoid tissue known as **Peyer's patches** which produce lymphocytes. The wall of the small intestine bears crypts between the base of villi called **crypts of Leiberkuhn** (Figure.5.3).

The large intestine consists of caecum, colon and rectum. The caecum is a small blind pouch like structure that opens into the colon and it possesses a narrow finger like tubular projection called **vermiform appendix**. Both caecum and vermiform appendix are large in herbivorous animal and act as an important site for cellulose digestion with the help of symbiotic bacteria. The colon is divided into four regions – an ascending, a transverse, a descending part and a sigmoid colon. The colon is lined by dilations called **haustra** (singular – haustrum) (Figure.5.4). The "S" shaped sigmoid colon (pelvic colon) opens into the rectum. Rectum is concerned with temporary storage of faeces. The rectum open out through the anus. The anus is guarded by two anal sphincter muscles. The anal mucosa is folded into several vertical folds and contains arteries and veins called anal columns. Anal column may get enlarged and causes **piles** or **haemorrhoids**.




Figure 5.3 Small intestine with Villi

TN_GOVT_XI_Bio Zoology_Ch05.indd 76 09-01-2020 5.27.55 PM