(iii) $\int \frac{1}{(3x-2)}dx = \frac{1}{3}\log|(3x-2)|+c$ (iv) $\int \frac{1}{(5-4x)}dx = -\frac{1}{4}\log|(5-4x)|+c$

**Example 11.7**

Integrate the following with respect to $x$:

(i) $\frac{1}{1+(2x)^2}$ (ii) $\frac{1}{\sqrt{1-(9x)^2}}$ (iii) $\frac{1}{\sqrt{1-25x^2}}$

**Solution**

(i) $\int \frac{1}{1+(2x)^2}dx = \frac{1}{2}\tan^{-1}(2x)+c$ (ii) $\int \frac{1}{\sqrt{1-(9x)^2}}dx = \frac{1}{9}\sin^{-1}(9x)+c$

(iii) $\int \frac{1}{\sqrt{1-25x^2}}dx = \int \frac{1}{\sqrt{1-(5x)^2}}dx = \frac{1}{5}\sin^{-1}(5x)+c$

## EXERCISE 11.2

Integrate the following functions with respect to $x$:

(1) (i) $(x+5)^6$ (ii) $\frac{1}{(2-3x)^4}$ (iii) $\sqrt{3x+2}$

(2) (i) $\sin 3x$ (ii) $\cos(5-11x)$ (iii) $\text{cosec}^2(5x-7)$

(3) (i) $e^{3x-6}$ (ii) $e^{8-7x}$ (iii) $\frac{1}{6-4x}$

(4) (i) $\sec^2\frac{x}{5}$ (ii) $\text{cosec}(5x+3)\cot(5x+3)$ (iii) $\sec(2-15x)\tan(2-15x)$

(5) (i) $\frac{1}{\sqrt{1-(4x)^2}}$ (ii) $\frac{1}{\sqrt{1-81x^2}}$ (iii) $\frac{1}{1+36x^2}$

## 11.5 Properties of Integrals

(1) If $k$ is any constant, then $\int kf(x)dx = k\int f(x)dx$

(2) $\int (f_1(x)\pm f_2(x))dx = \int f_1(x)dx \pm \int f_2(x)dx$

BDL7WU

**Note 11.1**

The above two properties can be combined and extended as

$$\int (k_1f_1(x)\pm k_2f_2(x)\pm k_3f_3(x)\pm\cdots\pm k_nf_n(x))\,dx$$

$$= k_1\int f_1(x)dx \pm k_2\int f_2(x)dx \pm k_3\int f_3(x)dx \pm\cdots\pm k_n\int f_n(x)dx.$$

That is, the integration of the linear combination of a finite number of functions is equal to the linear combination of their integrals

**Example 11.8**

Integrate the following with respect to $x$:

(i) $5x^4$ (ii) $5x^2-4+\frac{7}{x}+\frac{2}{\sqrt{x}}$ (iii) $2\cos x-4\sin x+5\sec^2 x+\text{cosec}^2 x$



11_11th-Maths-Vol2-Unit11.indd 190 23-03-2019 10:14:34