(5) Find the area of the parallelogram whose two adjacent sides are determined by the vectors $\hat{i}+2\hat{j}+3\hat{k}$ and $3\hat{i}-2\hat{j}+\hat{k}$.

(6) Find the area of the triangle whose vertices are $A(3,-1,2)$, $B(1,-1,-3)$ and $C(4,-3,1)$.

(7) If $\vec{a},\vec{b},\vec{c}$ are position vectors of the vertices *A, B, C* of a triangle *ABC*, show that the area of the triangle *ABC* is $\frac{1}{2}|\vec{a}\times\vec{b}+\vec{b}\times\vec{c}+\vec{c}\times\vec{a}|$. Also deduce the condition for collinearity of the points *A, B,* and *C*.

(8) For any vector $\vec{a}$ prove that $|\vec{a}\times\hat{i}|^2+|\vec{a}\times\hat{j}|^2+|\vec{a}\times\hat{k}|^2=2|\vec{a}|^2$.

(9) Let $\vec{a},\vec{b},\vec{c}$ be unit vectors such that $\vec{a}\cdot\vec{b}=\vec{a}\cdot\vec{c}=0$ and the angle between $\vec{b}$ and $\vec{c}$ is $\frac{\pi}{3}$. Prove that $\vec{a}=\pm\frac{2}{\sqrt{3}}(\vec{b}\times\vec{c})$.

(10) Find the angle between the vectors $2\hat{i}+\hat{j}-\hat{k}$ and $\hat{i}+2\hat{j}+\hat{k}$ using vector product.

BAZDFD

EXERCISE 8.5

Choose the correct or the most suitable answer from the given four alternatives

(1) The value of $\overrightarrow{AB}+\overrightarrow{BC}+\overrightarrow{DA}+\overrightarrow{CD}$ is

(1) $\overrightarrow{AD}$ (2) $\overrightarrow{CA}$ (3) $\vec{0}$ (4) $-\overrightarrow{AD}$

(2) If $\vec{a}+2\vec{b}$ and $3\vec{a}+m\vec{b}$ are parallel, then the value of *m* is

(1) 3 (2) $\frac{1}{3}$ (3) 6 (4) $\frac{1}{6}$

(3) The unit vector parallel to the resultant of the vectors $\hat{i}+\hat{j}-\hat{k}$ and $\hat{i}-2\hat{j}+\hat{k}$ is

(1) $\frac{\hat{i}-\hat{j}+\hat{k}}{\sqrt{5}}$ (2) $\frac{2\hat{i}+\hat{j}}{\sqrt{5}}$ (3) $\frac{2\hat{i}-\hat{j}+\hat{k}}{\sqrt{5}}$ (4) $\frac{2\hat{i}-\hat{j}}{\sqrt{5}}$

(4) A vector $\overrightarrow{OP}$ makes 60° and 45° with the positive direction of the *x* and *y* axes respectively. Then the angle between $\overrightarrow{OP}$ and the *z*-axis is

(1) 45° (2) 60° (3) 90° (4) 30°

(5) If $\overrightarrow{BA}=3\hat{i}+2\hat{j}+\hat{k}$ and the position vector of *B* is $\hat{i}+3\hat{j}-\hat{k}$, then the position vector *A* is

(1) $4\hat{i}+2\hat{j}+\hat{k}$ (2) $4\hat{i}+5\hat{j}$ (3) $4\hat{i}$ (4) $-4\hat{i}$

(6) A vector makes equal angle with the positive direction of the coordinate axes. Then each angle is equal to

(1) $\cos^{-1}\left(\frac{1}{3}\right)$ (2) $\cos^{-1}\left(\frac{2}{3}\right)$ (3) $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$ (4) $\cos^{-1}\left(\frac{2}{\sqrt{3}}\right)$

08_11th-Maths-Vol2-Unit8.indd 80 23-03-2019 10:02:28