4.7.5 Positions used for patients.

Positions

RYVBDF

1. Dorsal position (Supine.):

Patient is flat on the bed with legs extended and arms at the sides of the body. This is not a comfortable position, as the curves of the body are not supported.




Indications

- Surgical procedures, it allows access to the peritoneal, thoracic and pericardial regions; as well as the head, neck and extremities.

2. Dorsal recumbent position:

Place patient flat on back with one pillow under head; have knees flexed and separated and feet flat on bed.



Indications

- Rectal, vaginal and pelvic examinations and treatments.
- Deliveries.

3. Lateral Position:

Patient lies on his side with spine straight. The knees are flexed; the upper knees are more flexed than the lower one. Pillows may be provided for the head, in between the legs, and to support the back and abdomen. The lower arm is kept above the head and the upper arm is placed on a pillow in front.



Indications

- General comfort, rest and relaxation.
- Back care.

The arms and legs do not bear the weight of the body.

4. Sims or Left Lateral position:

Place patient on left side somewhat obliquely across the bed with buttocks to edge of mattress. Incline the body forward, draw the left arm back under patient and place the right arm free in front. The thighs should be flexed upon the body, the right more than the left.



Indications

- Vaginal examinations.
- Perineal examination.
- Rectal examinations.
- Post operative, to maintain a clear airway.

5. Jack knife position:

Place patient on a prone position with the hips directly over the band of the examining table. Tip the table with the head lower than

Nursing-voc_Unit 04.indd 103 22-02-2019 4.44.52 PM