the hips. Lower the foot part of the table so that the patient's feet are below the level of his head.



Indications

- For drainage after any procedures.
- Operation on the rectum and coccyx.

6. Knee Chest Position:

Place patient in the prone position, then assist her to kneel so that her weight rests on her chest and knees. Turn head to one side and flex her arms at the elbows extending, then to the bed in front of her. Be sure the thighs are perpendicular to the level of the head. Watch pulse and general condition of the patient.



Indications

- To obtain better exposure of the vagina, cervix, and rectum.
- To examine the bladder.
- To help correct retroversion of the uterus.
- To administer caudal and sacral anesthesia.
- Vaginal and rectal examinations.
- Operative procedures on the vagina, rectum and perineum.
- Operative deliveries

7. Lithotomy Position:

A position of the body for medical examination, pelvic or abdominal surgery, or childbirth in which the individual lies on the back with the hips and knees flexed and the legs spread and raised above the hips often with the use of stirrups.



Indications

- Abdominal surgeries.
- Childbirth.
- Pelvic examination.
- Urologic examination of the prostate.
- Male urethral surgery. Examination or operations on rectum and genital organs.

8. Prone Position:

Patient lies flat on his abdomen with head kept on a pillow and turned to one side and another pillow under the lower chest. Pillows are kept under the waist and under the lower legs. The arms are flexed at the elbow and kept above the head.



Nursing-voc_Unit 04.indd 104 22-02-2019 4.44.52 PM