Highlight

Strongest nurse stands beside patient's head &shoulders, or beside female patient's hips.

STUDENT'S ACTIVITY

Demonstration of moving, shifting and lifting methods on each other.

Fact.

In one recent study, people who took ***12 weeks of yoga classes*** had fewer symptoms of low back pain.

4.12 OXYGEN NEEDS.

4.12.1 Introduction

Oxygen is administered whenever there is deficiency in the blood and is shown by cyanosis. Patients with respiratory dysfunction are treated with oxygen therapy to relieve anoxaemia or Hypoxemia. The normal amount of oxygen in the blood must be in the range of 80 to 100 mm of Hg. If it falls below 60 mm of Hg, irreversible physiological effect may take place. The brain cells receive 20 percent of the body's oxygen supply and can live only for 3 to 5 minutes if the oxygen supply is cut off.

4.12.2 Purpose

- To supply oxygen in conditions when there is interference with the normal oxygenation of the blood.

4.12.3 Indications for Oxygen Inhalation

1. Breathlessness due to asthma, pulmonary embolism, emphysema, cardiac insufficiencies etc.
2. Obstructed airway due to growth, and enlarged thyroid.
3. Cyanosis.
4. Shock and circulatory failure.
5. After severe haemorrhage.
6. Anaemia.
7. Patients under anaesthesia.
8. Asphyxia due to any reason e.g. drowning, inhalation of poisonous gases, hanging etc.
9. Poisoning with chemicals that alter the tissues ability to utilize oxygen e.g. cyanide poisoning
10. Carbon monoxide poisoning.
11. Postoperative chest surgery and thyroidectomies.
12. Insufficient oxygen in the atmosphere.
13. Air hunger.

4.12.4 Methods of Oxygen Administration.

Administration of oxygen depends upon the condition of the patient, the concentration desired, the facility available and the preference of the doctor. It can be given by the following ways:

Methods

1. Oxygen by nasal catheter/cannula:

This is the usual method of administering oxygen to the patients in the ward. The nasal

Nursing-voc_Unit 04.indd 120 22-02-2019 4.44.57 PM