catheter is inserted into the nostril reaching upto the uvula. The catheter is taped on the forehead to keep it in place. The nasal catheter permits free movement for the patient and nursing care may be given with much more ease.



2. Nasal prongs:

This is another method of administering oxygen to the patients in the ward.



3. Oxygen by B. L. B. Mask:

When oxygen concentration of over 25% is needed or when oxygen is given under pressure, the mask is used. If the mask does not fit snugly over the face, oxygen will be lost from the mask. It is useful for the patients who are unable to breathe through the nose. Flow of 8 to 12 litres oxygen will be sufficient to maintain the concentration of oxygen from 25 to 60%. B. L. B. [Boothby, Lovelace and Bulbalian] is a rubber mask. It is made to fit over the nose or nose and mouth. The reservoir bag is attached to it.



4. Oxygen by Tent:

It consists of a canopy over the patient which may cover the patient partially or totally. It is connected to a supply of oxygen. The canopies are transparent, so that the nurse can observe the patient. The lower portion of the canopy is tucked under the bed to prevent the escape of oxygen.



4 (a) oxygen hood:

This is used for infants to administer oxygen

5. Oxygen cylinders and accessories:

Oxygen is supplied in cylinders or tanks. It is stored under a pressure of 2200 lbs /1000 kg per sq. inch. Oxygen is stored in the oxygen cylinder are low pressure about 50 to 60 lbs per sq. inch. The oxygen cylinders are painted black with white neck. The Wolfs bottle has two holed rubber cork in which two glass tubes are inserted one short and one long. The long tube is attached by rubber tubing to the oxygen cylinder. The short tube is attached by the rubber tubing

Nursing-voc_Unit 04.indd 121 22-02-2019 4.44.59 PM