symmetric expansion and the thumbs separate 2–3 cm during deep inspiration when assessing for the respiratory excursion. The client manifested quiet, rhythmic and effortless respirations.

Breast: Whether there are any abnormalities in the shape and size. Whether there are any lumps or discharge from the nipples.

Abdomen: The abdomen of the client has an unblemished skin and is uniform in color. The abdomen has a symmetric contour. There were symmetric movements caused associated with client's respiration.

Upper Extremities: The extremities are symmetrical in size and length.

Muscles: The muscles are not palpable with the absence of tremors. They are normally firm and showed smooth, coordinated movements.

Bones: There were no presence of bone deformities, tenderness and swelling.

Joints: There were no swelling, tenderness and joints move smoothly. Whether the nails are broken or brittle. Whether there is clubbing of fingertips, tremors of hands, swelling of extremities, pain in the joints or any other abnormality, all range of motions present

6.6 PHYSIOLOGICAL ASSESSMENT

The most frequently measurement obtained by health care providers are those of temperature, pulse, blood pressure and respiration as the indicators of health status, these measures indicate the effectiveness of circulatory, respiratory, neural and endocrine body functions.

Vital signs include the physiological measurements of temperature, pulse, Bp and respirations. Vital signs are a quick and efficient way of monitoring a patient's response to intervening changes. One vital sign can influence characteristics of other vital signs.

Assessment of vital signs allows the nurse to identify nursing diagnoses, to implement planned intervention and to evaluate success. When the nurse learns the physiological variables influencing vital signs and recognizes the relationship of vital sign changes to other physiological assessment findings.

Vital Sign Ranges			
Age-Appropriate Vital Signs			
	Heart Rate	Respirations	Blood Pressure
Newborn	80–180	30–60	60–80/30–60
Toddler	80–110	24–32	90–100/50–65
School Age	60–110	18–26	95–110/55–70
Adolescent	50–90	16–20	110–120/60–80
Adult	60–100	12–20	110–140/60–90

Nursing-voc_Unit 06.indd 176 22-02-2019 4.45.55 PM