early morning gradually arises during morning and afternoon and peaks in evening.

- **Gender:** After puberty males tend to have higher BP. After, menopause, women tend to have high BP. BP is measured by sphygmomanometer.

## Purposes of Monitoring BP

1. To aid in the diagnosis of the patient's condition.
2. To guide in his treatment.
3. To evaluate the patient's progress.

## General instructions

1. See that the patient is relaxed and is a comfortable position.
2. Help to take blood pressure for patients with the following conditions:
   - New patients.
   - Pre and post-operative patients.
   - Antenatal and post-natal patients.
   - Patients with shock and hemorrhage.
   - Patients with cardiac conditions and hypertension.
   - Patients with neurological disorders.
3. Record pulse along with blood pressure.
4. Blood pressure is taken at the same arm, same time, and same posture daily.

## Equipments

1. Sphygmomanometer
2. Stethoscope
3. Pen



**Sphygmomanometer**




Do you know who invented the BP apparatus?
The sphygmomanometer was invented by Samuel Siegfried Karl Ritter Von Basch in 1881.

## Guidelines for taking BP

- The sphygmomanometers generally used in clinical setting are mercury type. And aneroid type. The mercury type sphygmomanometers are more reliable than the aneroid type sphygmomanometers. The aneroid sphygmomanometer gives blood pressure reading on dial indicator.
- Systolic pressure is increased in pressure induced by systolic contraction and diastolic pressure is decrease in pressure induced by diastolic relaxation of the left ventricle of heart.
- Never take blood pressure when the patient is excited, exhausted and just after exercise, smoking or meals.
- Allow the patients to rest for five minutes before taking blood pressure.



Nursing-voc_Unit 06.indd 189 22-02-2019 4.45.58 PM