comfortable position. Support the arm.

- Expose the arm and keep it extended.
- Apply the end of the cuff with the rubber bag over the brachial artery two inches above the elbow.
- Apply the end of the cuff smoothly and snugly around the upper arm. Tuck the end neatly.
- Place the sphygmomanometer in position.
- Stay with the patient until the procedure is over.
- Remove the cuff from the patient's arm, roll neatly and replace in the box. See patient is comfortable.
- The reading is recorded in the chart.
- Systolic pressure is always written over the diastolic pressure E.g. 80/120mmHg.




Recording of Blood Pressure

Variation in BP


Hypertension: Elevated or high blood pressure is known as hyper tension. Hyper tension is a major factor causing deaths from strokes and myocardial infarction (Heart arrest)

Hypotension: When the systolic pressure falls to 90 mm Hg or below, that condition is known as hypotension.



6.6.5 Pain

It is a distressing feeling often caused by intense or damaging stimuli. Because it is a complex, subjective phenomenon, defining pain has been a challenge.



STUDENT'S ACTIVITY

A 42-year-old woman has complained the severe pain in the lower quadrant of the abdomen for 6 hours. She has had nausea and vomiting for two days. She was pale and painful. How do you assess the patient's condition?

Nursing-voc_Unit 06.indd 191 22-02-2019 4.45.58 PM