Closed fracture	The surrounding skin is unbroken but internal injury to surrounding tissue.

Signs and symptoms

- Difficulty in moving a limb normally.
- Pain at or near the site of injury, made worse by movement.
- Tenderness over a bone if gently touch is a sign of fracture.
- Distortion, swelling and brushing at the site of the fracture.
- Coarse grating of the bone end may be heard or felt.
- A shortening, bending, or twisting of the affected limb.

To Remember

- DO NOT move the casualty until the injured part is secured and supported unless he/she is in danger
- DO NOT let the casualty eat or drink
- DO NOT try to replace a disclosed bone into its socket

TREATMENT



For open fracture

a. Cover the wound with clean pad or sterile dressing, apply pressure to control the bleeding.
b. Without touching an open wound with your fingers, carefully place some clean padding over and around the dressing.
c. Secure the dressing and padding. Bandage firmly, but not tightly that the circulation is impeded.
d. Immobilize the injured part as for a closed fracture

For the closed fracture

a. Tell the casualty to keep still, steady and support the injured part with your hands until it is immobilized.
b. For firmer support, secure the injured part to a sound part of the body. And bandage from the uninjured side.



STUDENT'S ACTIVITY

On a walk through your neighborhood with friends, you find a man lying on the ground under a ladder. He is in obvious pain and his arm is clearly broken, with a piece of bone protruding from the skin. How to handle the situation?

c. Check the circulation beyond any bandages every minute.

Nursing-voc_Unit 07.indd 205 2/23/2019 2:50:14 PM