Signs:-

- Pulse rate increased
- Blood pressure falls
- Pupils are dilated
- Lustreless eyes
- Shaking and trembling of arms and legs
- Unconsciousness may develop.

Types of Shock	
Neurogenic	Spinal or head injury resulting in loss of nerve control
Haemorrhagic	Loss of blood due to wound and internal bleeding.
Respiratory	There is an insufficient amount in the blood due to inadequate breathing
Cardiac	Cardiac muscle not pumping effectively due to heart attack
Metabolic	Loss of body fluids with a change in biochemical equlibrium
Septic	Severe infection can cause septic shock
Anaphylactic	Severe allergic reaction of the body to sensitization by a foreign protein

Management


a. Immediately reassure and comfort the casualty.

b. Body positioning for shock.



- DO NOT let the casualty smoke, eat drink, or move unnecessarily. If he complaints of thirst moisten her lips with little water.
- DO NOT leave the casualty unattended.
- DONOT try to warm casualty with a hot water bottle or any other direct source of heat.

c. Normally the lower extremities should be elevated. By gravity this reduce the blood in the extremities and may improve the blood supply to the heart.



d. If there are indications of the head injuries, the head could be raised slightly to reduce pressure on the brain.

e. If there are breathing difficulties, the victim may be more comfortable with head and shoulders raised

f. Loosen the tight clothing to help the circulation and assist breathing.

g. Treats the cause of shock, stop bleeding, immobilize fracture.

h. f breathing and heart beat stop then;-

i. Establish the airway

- Begin resuscitation immediately.
- Keep patient in recovery position.

Nursing-voc_Unit 07.indd 207 2/23/2019 2:50:16 PM