5. **Seek medical assistance:**
   - If the wound appears to be minor and the patient is able to travel by car, arrange an urgent appointment with a local doctor to assess and treat the injury.
   - If the injury is severe or the patient is very unwell – call 108 for an ambulance as soon as possible.
   - While waiting for an ambulance to arrive, observe the patient closely for any change in condition.
6. **If blood leaks through the pressure pad and bandage:**
   - Apply a second pad over the first. Use a tea towel or similar bulky fabric and apply maximum pressure to the area.
   - For major uncontrolled bleeding quickly remove the blood-soaked pad and bandage then replace with a fresh bulky pad and bandage. The continuing bleeding may be due to the pad slipping out of position when the first bandage was applied.



NOSE BLEED

A blow to the nose, flying at high altitude, or scuba diving may all cause a bleeding nose. For a child, always check whether there is a foreign body present – e.g. a bead or coin. If this has occurred, seek prompt medical advice and DO NOT try to remove the object yourself because this may cause further damage.



Management

1. Sit the casualty down with her head held well forward.
2. Ask the patient to breath through her mouth and to punch her nose just below the bridge. Help her if necessary.



> ***Caution.....?***
> DO NOT let her head back, blood may run down her throat and induce vomiting

3. Tell her not to speak, swallow, cough, spit or shift.
4. Give her clean cloth or tissue to mop her dribble.
5. After ten minutes, tell the casualty to release the pressure. If her nose is still bleeding reapply the pressure for the further periods of ten minutes.
6. Once the bleeding is under control and with the casualty still leaning forward

Nursing-voc_Unit 07.indd 213 2/23/2019 2:50:19 PM