- To use the head – tilt/chin – lift method, place the palm of your hand that is closest to the victim's head on forehead and your other hand under his neck. Place the hand lifting his neck close to the back of his head to minimize cervical-spine extension.



**Air way**

- Then gently press back on his forehead while lifting up and supporting his neck.
- Use the jaw – thrust without head – tilt method if you suspect the victim has a neck or spine injury. Kneel at the victim's head, facing his feet.
- Listen for any air movement and look to see if his chest or abdomen is moving up and down. Feel with your chest for any flow of air. If the victim has started to breathe, maintain airway until help arrives.

## Giving mouth to mouth respiration

a. With the casualty lying flat on his back remove any obvious obstruction., including broken or displaced dentures from the mouth.

b. Open the airway by tilting the head and using two fingers to lift the chin.

c. Close the casualtynose by pinching it with your index finger and thumb. Take full breath, and place your lips around his mouth making a good seal.

d. Blow into the casualty mouth until you see the chest rise. take about two second for full inflation.



## Mouth -To – mouth Respiration

e. Remove your lips and allow the chest to fall fully, which takes four seconds. Repeat this once and then assess for the sign of circulation.

f. When you see the victim's chest rise, then fall, you will know that air is entering and escaping his lungs.

## Giving chest compression

a. Check the victim's carotid pulse for ten seconds.

b. Kneel bedside the casualty. locate one 4th and 5$^{th}$ intercoastal space

c. Place the heel of your hand on the breast bone That is the where you should apply pressure.

d. Place heel of your first hand on top of the other hand, and interlock your fingers.



Nursing-voc_Unit 07.indd 222 2/23/2019 2:50:23 PM