e. Leaning well over the casualty with your arms straight, press vertically down and depress the breastbone approximately–5-4 cm. Release the pressure without removing hands.

f. Compress the chest 15 minutes aiming for a rate of 100 compression per minute. Then give two breaths of artificial ventilation.

g. If you are only rescuer, time your compressions at a rate of 100 a minute. Count, "one two and three and four and five and ...." Up to the count of fifteen.

h. Then deliver two quick breaths without allowing the victim to exhale between them.

i. Perform CPR for 1 minute; check the victim's pulse then quickly telephone for help if none has arrived. Return quickly and resume CPR. If there is no phone available, continue CPR.

Cardiopulmonary resuscitation for small children

a. Use adult CPR techniques for children older than 8 years. In emergency, of course, you are not going to delay CPR until you determine the child's age. Instead, consider his body size relation to the size of your hand.



Cardiopulmonary Resuscitation

b. For example, if looks too small to use both hands for cardiac compression, use the heel of one hand. If he is too small for that, use two, three fingers.

c. If he has a small face, place your mouth over his mouth and nose, when ventilating, give only enough air to make the child's chest rise.

d. Try to palpate the child's carotid pulse. If you find a pulse. Do not give cardiac compression but do ventilate the child a rate of one breath every 4 seconds.

e. If you can't locate a pulse, find the proper location for compression. Use the same technique you would for an adult.

f. Then compress about 1 to 1. ½ inches (2.5 to 3.8 cm), using the heel of one hand (as shown).

7.20 BANDAGES AND SLINGS

Bandages have a number of purposesthey are used to hold dressing in position over wounds, to control bleeding, to support and immobilize injuries, and reduce the swelling.

General Rules for bandaging

a. Reassure the casualty and explain clearly what you are going to do.

b. Bandages should be applied firm enough to keep dressing and splints in position.

c. But no so tight as to cause injury to the part or to impede the circulation of the blood.

d. A bluish tinge of the finger or nails may be a danger sign that the bandages are too tight.

e. Loss off sensation is another sign.

Nursing-voc_Unit 07.indd 223 2/23/2019 2:50:23 PM