Ear bandage

Lay the outer surface of the bandage against forehead and carry the bandage round the head in one circular turn, bandaging away from the injured ear. Towards the sound side, carry the bandage round to the back of the head, low down in the nape of the neck again, repeat these.



Ear bandage

Eye bandage

Lay the outer surface of the bandage against the forehead and take the circular turn round the head, bandaging away from the injured eye. Carry the bandage on, round side for the second time. Take it obliquely to the back of the head, under the prominence at the back of the skull and from there bring it upwards beneath the ear of the affected side, over the pad of the circular turn and continue.



Eye bandage

Tubular gauze bandage

This is a special form of tubular bandage, which can be applied with an applicator to any part of the body, it is ideal for small dressing on hands and limbs.

1. Inspect the nails to see, if there is any bluish colour. A bluish colour shows that there is a dangerous tightening of splint or plasters and therefore, free flow of blood is not possible.
2. If the casualty is not wearing a coat, place a soft pad under the neck portion of the sling to prevent rubbing of the skin in that place.

Collar and cuff sling

This sling is used to support the wrist only.

1. The elbow is bent and the forearm is placed across the chest in such a way that the fingers touch the opposite shoulder. In this position, the sling is applied.
2. A clove hitch is passed round the wrist and the ends tied in the hollow above the collar bone on the injured side.



Collar and cuff sling

Nursing-voc_Unit 07.indd 227 2/23/2019 2:50:27 PM