Procedure:

1. Place all the articles at conveniently on the bed side table.
2. Explain the procedure to the patient.
3. Put the rubber sheet (mackintosh) with towel and kidney tray under the chin.
4. Have the patient rinsed his mouth with salt solution from the feeding cup.
5. Turn the patient's head to one side.
6. Take the artery forceps, wrap a piece of rag pieces/cotton ballaround the tip of the forceps.
7. Dip inside the saline water and clean the teeth with up and down movements.
8. Pay special attention to inside the mouth.
9. Change rag pieces or cotton balls as often if necessary.
10. Discard used cotton in the other kidney tray.
11. Allow the patient to gargle as much as necessary.

Care of Dentures

If the patient has dentures, care should taken to keep the dentures clean. If the patient is unable to do so, the nurse has to remove the dentures by grasping it with gauze pieces. Place them in a tumbler or cup containing water. Dentures are washed carefully using brush, toothpaste and cool water. Water, which is too hot, may injure the composition of dentures. If the patient is to do by himself, he may be assisted. Remove dentures of patients who are unconscious, mentally ill and who have vomiting, cough or spasm.

12. Dip the swab stick in glycerin borax, swab gums, root and sides of the mouth.

After care equipments:

1. Clean kidney trays and feeding cups with soap and water.
2. Boil forceps and gallicups after cleaning
3. Place all articles in their places after cleaning and boiling.

2.3 EYE CARE:

A common problem of the eyes are secretions that dry on the lashes as crusts. If new borns, the eyes are treated soon after the baby is born to prevent opthalmia neonatorum. Eyes are cleaned from the inner to the canthus.

Articles Required:

1. Mackintosh and towel.
2. Sterile bowl with sterile cotton swabs.
3. Sterile normal saline or any other ordered solution.
4. Kidney tray and paper bag.
5. Clean face towel.

Procedure:

1. Wash hands.
2. Pour sterile saline into the bowl and wet the cotton swabs.
3. Stand infront of the patient clean the eyes with the sterile swabs. Discard the swabs into the paper bag continue cleaning till all discharges are removed from the eyes.
4. For crusted secretion, place a wet warm gauze piece or cotton swab over the closed eye. Leave it in place until the crust becomes soft.

Nursing-Voc_Practical_Unit 1-5.indd 272 23-02-2019 3.17.28 PM