

12. Fowler's Position and Semi-Fowler's Position:


Patient is in a partially sitting position. The back of the bed is elevated to 45 degrees with the aid of a backrest and pillow or by adjustment of the cot. It can be elevated to 30 degrees as well as 90 degrees. Patient's back shoulder and head are supported well. The knees are flexed and supported with a pillow or by cot adjustment. A footrest is provided to prevent foot drop.



WT30⁰



UT60⁰




WT60⁰

Indications

- To obtain good drainage in the pelvis.
- To localize infection in the pelvis and prevent it's spread to the peritoneum.
- To prevent strain of abdominal muscles.
- This position is used for patients with dyspnoea (difficulty in breathing), distended abdomen, abdominal surgery, cardiothoracic disorders and ascites.
- The position is also useful while passing Ryle's tube. And while performing tapping of ascites fluid.

13. Trendelenburg position:

The patient lies on his back with the foot at the bed elevated on wooden blocks. Patient's head and trunk are lower than the legs.



Reverse Trendelenburg Position:

The head and shoulders are at a higher level than the hips, legs and feet. This position is used for reducing intracranial pressure and for other treatment measure.



Indications

- Gynecological surgery and suprapubic prostatectomy cases.
- To prevent shocks.
- To prevent or relieve post-partum hemorrhage.

Nursing-Voc_Practical_Unit 1-5.indd 289 23-02-2019 3.17.35 PM