- one end being continued round the scalp and other going order it
- scalp turn secured by horizontal turn
- capline bandage completed

Eye and Ear Bandage





10.4 PATTERNS USED IN BANDAGING

1. Circular turns, as used for head and trunk.
2. Simple spiral, for parts of uniform thickness, eg. Fingers wrist.
3. Reverse spiral, used on limbs where the thickness of the part varies, e.g forearm leg.
4. Figure-of-Eight

This may be used on limbs instead of the reverse spiral also for the hand and foot.

5. Spica, used for the shoulder, hip and thumb
6. Divergent Spica, for a flexed joint, e.g elbow, knee, heel
7. Recurrent to cover tips of fingers or a stump.
8. Special bandages such as the capeline for the head, eye bandage, ear and breast bandages.

10.5 APPLICATION OF BANDAGE

Preliminary Assessment

- Check the doctors order to see the specific precautions if any regarding the positioning and movement.
- Assess the patients need for application of bandage.
- Monitor vital signs.
- Assess the patients mental status.
- Assess the need for pain medication
- Assure the patient, the patient's family.
- Assess the adequacy of circulation by noting surface temperature, skin colour, and sensation of body parts to be wrapped.
- For tying the bandage a 'reef knot' must be always used.
- The knot should be made where it does not hurt the skin or cause discomfort.
- Tuck the loose ends of the bandage out of sight.
- Not in use the triangular bandages should be folded narrow. Bring the two ends to the centre and fold again. It becomes a packet which measures 16 x 9 cm handy to carry.
- Wrinkled Bandages are uncomfortable.
- Never ignore any complaints of pain experienced by the patient. This should be invested and the cause is removed immediately.
- Do not use extra turns in order to use all the bandages.

Nursing-Voc_Practical_Unit 10.indd 308 23-02-2019 4.06.25 PM